SAM SILVERMAN LISA A. TELL

RADIOLOGY OF BIRDS

An Atlas of Normal Anatomy and Positioning















SAUNDERS

RADIOLOGY OF BIRDS

An Atlas of Normal Anatomy and Positioning

RADIOLOGY OF BIRDS

An Atlas of Normal Anatomy and Positioning

SAM SILVERMAN, DVM, PHD, DIPL ACVR

Clinical Professor

Department of Surgical and Radiological Sciences
School of Veterinary Medicine
University of California, Davis
Davis, California

LISA A. TELL, DVM, DIPL ABVP (AVIAN), ACZM

Professor

Department of Medicine and Epidemiology
School of Veterinary Medicine
University of California, Davis
Davis, California

RADIOGRAPHIC and TECHNICAL ASSISTANCE JODY NUGENT-DEAL and KRISTINA PALMER-HOLTRY

Registered Veterinary Technicians
Companion Avian and Exotic Pet Medicine Service
Veterinary Medical Teaching Hospital
University of California, Davis
Davis, California

ARTISTIC ILLUSTRATIONS

KATHY WEST, M.SC. Scientific and Medical Illustrator Davis, California

with 485 illustrations





11830 Westline Industrial Drive St. Louis, Missouri 63146

RADIOLOGY OF BIRDS

ISBN-13: 978-0-7216-0635-4 ISBN-10: 0-7216-0635-0

Copyright © 2010 by Saunders, an imprint of Elsevier Inc.

All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the publisher. Permissions may be sought directly from Elsevier's Rights Department: phone: (+1) 215 239 3804 (US) or (+44) 1865 843830 (UK); fax: (+44) 1865 853333; e-mail: healthpermissions@elsevier.com. You may also complete your request on-line via the Elsevier website at http://www.elsevier.com/permissions

Notice

Neither the Publisher nor the [Authors] assume any responsibility for any loss or injury and/or damage to persons or property arising out of or related to any use of the material contained in this book. It is the responsibility of the treating practitioner, relying on independent expertise and knowledge of the patient, to determine the best treatment and method of application for the patient.

The Publisher

Library of Congress Cataloging-in-Publication Data

Silverman, Sam

Radiology of birds : an atlas of normal anatomy and positioning / Sam Silverman, Lisa A. Tell ; radiographic and technical assistance, Jody Nugent-Deal and Kristina Palmer-Holtry ; artistic illustrations, Kathy West.

p.; cm.
Includes index.
ISBN-13: 978-0-7216-0635-4 (hardcover: alk. paper)
ISBN-10: 0-7216-0635-0 (hardcover: alk. paper)
ISBN-10: 1. Birds--Anatomy--Atlases. 2. Veterinary radiography--Atlases. I. Tell, Lisa A. II. Title.
[DNLM: 1. Birds--anatomy & histology. 2. Radiography--veterinary. QL 697 S587r 2010]
SF767.B57S55 2010
636.6'08960757--dc22

2009008520

Vice President and Publisher: Linda Duncan Senior Acquisitions Editor: Anthony Winkel Associate Developmental Editor: Lauren Harms Publishing Services Manager: Julie Eddy Senior Project Manager: Celeste Clingan Design Direction: Kim Denando

Working together to grow libraries in developing countries

www.elsevier.com | www.bookaid.org | www.sabre.org

ELSEVIER

BOOK AID

Sabre Foundation

PREFACE

The species included in this text are either popular companion pets or birds that are commonly presented to rehabilitation facilities. Documentation of radiographic and alternative imaging findings associated with disease conditions of these species continues to expand. There are several other text-book references documenting normal radiographic imaging anatomy for birds; however, this text was designed to provide additional radiographic information regarding normal anatomy and provide examples of contrast radiographic studies and advanced imaging procedures.

Given the limitations in printing, reproducing radiographic images of animal subjects such as birds is a challenge. It is difficult to obtain print quality that will allow the reader to appreciate subtle changes in the contrast and detail while still making the text affordable. Therefore, images in this text may not fully represent the image quality of the original radiographs for some avian species. In order to supplement the printed version of the text, a companion CD containing the images has been included and the images on this format medium are a closer approximation of the image quality of the original studies.

The purpose of this atlas is to provide veterinarians with normal radiographic images, contrast studies, and alternative imaging studies. We hope this atlas increases the utilization and accuracy of diagnostic imaging of these avian species and enhances their medical care.

DEDICATION

This book is dedicated to the veterinarians, technicians, and students who have helped to advance the medical care of our feathered companions.

Sam Silverman and Lisa Tell

This text and its companion volume, *Radiology of Rodents*, *Rabbits, and Ferrets*, are the product of 7 years of effort by the authors. I could not ask for a harder working, more conscientious, and organized co-author than Dr. Lisa Tell.

The support and encouragement of my mentors must be recognized. Dr. James Ticer, the first radiologist to enter full-time private radiology practice, encouraged me to pursue radiology as a specialty and provided me the opportunity to author a chapter in his textbook, *Radiographic Technique in Small Animal Practice*. To the best of my knowledge, this is the first inclusion of an avian radiology chapter in a veterinary radiology textbook. My career has been shaped by many, but Drs. Joe P. Morgan, Peter F. Suter, and Timothy R. O'Brien provided the best veterinary radiology training imaginable, encouraged me to apply my skills to the imaging of avian patients, and stressed the importance the discipline and the scientific method. The knowledge and techniques I have gained through the years are a product of their dedication to teaching.

There were many long days, weekends, and evenings involved in the production of these texts and they were never questioned by my family. I acknowledge the sacrifices made by my wife, Debrah Tom, and children, Naomi, Emily, and Darrell

The journey that I embarked upon in the creation of Volumes 1 and 2 of these radiology texts was initiated by Dr. Sam Silverman. For giving me this opportunity and for his ability to remind me of the amazing gifts of life and health, I will always be grateful. Kathy West was an incredible person to work with and I especially appreciated her balanced approach to life. In the process of creating this book, I have gained lifelong friends in both Sam and Kathy.

My career in veterinary medicine has been deeply enriched by several of my mentors for whom I will forever be appreciative for all they have done for me. My interest in birds was sparked by my high school biology teacher, Mr. Larry Thorngren, who taught me to look in the sky and enjoy the feathered animals with which we share our world. Drs. Mitchell Bush and Scott Citino are amazing veterinarians; their dedication to teaching and passion for veterinary medicine were the basis for my training in the true "art" of zoological veterinary medicine. Lastly, I would like to thank Dr. Bill Lasley for providing endless encouragement and support and for mentoring me through my academic career

I would also like to dedicate this book to my parents, William and Bette Tell, and sister, Lee Ann Hughes, for all of their love, support, and inspiration. Their encouragement has allowed me to follow my passion for veterinary medicine. My life would not be complete without my husband, Don Preisler, and children, Nicholas and Alexander Preisler, and mother-in-law, Dawn Preisler, as they are a constant source of love and happiness.

Lisa Tell

Sam Silverman

ACKNOWLEDGMENTS

The untiring and talented efforts of many individuals made this atlas a reality. Hundreds of radiographic images were produced by Jody Nugent-Deal and Kristina Palmer-Holtry. Candi Stafford and Michelle Santoro provided technical advice and assistance to optimize image quality and develop radiographic protocols. Bob Smith was our technical liaison with the 3M Corporation. Jason Peters and Richard Larson developed the protocols for the computed tomography and magnetic resonance examinations. Dr. Allison Zwingenberger was exceptionally helpful evaluating the computed tomography and magnetic resonance studies. John Doval provided technical advice on the reproduction of images. Jacki Pessenti and Brett Stedman were responsible for the care of some of the birds used for imaging. Debrah Tom paid exceptional attention to detail required in developing the keys for the text. Kathy West's artistic, organizational, and scientific skills are responsible for the high quality of the text's medical illustrations. The fortitude, resourcefulness, and attention to detail of the aforementioned individuals resulted in production of quality images that serve as the basis of this text.

The animals imaged were integral to this text's creation. They were housed and maintained in compliance with the Animal Welfare Act and the Guidelines for the Care and Use of Laboratory Animals. All radiographic and alternative imaging procedures were performed according to an approved animal care and use committee protocol. We are indebted to the individuals who cared for these animals and provided permanent homes for them when imaging was completed.

The concept of this atlas evolved because of the vision of Ray Kersey, former Executive Editor of Veterinary Medicine for Elsevier. His dedication to disseminating veterinary medical knowledge cannot be understated.

Dr. Anthony Winkel, Linda McKinley, Maureen Slaten, and Lauren Harms of Elsevier provided invaluable assistance during the development of this textbook and at the final production phase of this atlas. Their dedication, hard work, and guidance regarding technical and esthetic matters were crucial and greatly appreciated.

Production of this atlas entailed four years of work and we would be remiss not to acknowledge the goodwill, support, and understanding of our friends and families. Our spouses, Don Preisler and Debrah Tom, constantly provided support and encouraged us to continue even though there were many hours devoted to the creation of the atlas rather than family affairs. Alexander and Nicki Preisler were good sports and were willing to lend their "Mom" to the spending time with the "Bird Book." Kathy West's husband, Ed West, was always willing to open his home to endless weekends and cooked delicious meals when we worked late into the night

In closing, we would like to extend our appreciation to all of these individuals for their encouragement, efforts, and sacrifices. We are indebted to them for their persistence and dedication.

Sam Silverman and Lisa Tell

CONTENTS

- 1 Radiology Equipment and Positioning Techniques, 1
- 2 Anatomic Art, 17
- 3 Budgerigar (Melopsittacus undulatus), 25
- 4 Peach-Faced Lovebird (Agapornis roseicollis), 29
- 5 Cockatiel (Nymphicus hollandicus), 39
- 6 Sun Conure (Aratinga solstitialis), 47
- 7 Senegal Parrot (Poicephalus senegalus), 51
- 8 African Grey Parrot (Psittacus erithacus erithacus), 67
- 9 Orange-Winged Amazon Parrot (Amazona amazonica), 75
- 10 Blue and Gold Macaw (Ara ararauna), 125
- 11 Goffin Cockatoo (Cacatua goffiniana), 147
- 12 Sulphur-Crested Cockatoo (Cacatua galerita), 151
- 13 Umbrella Cockatoo (Cacatua alba), 155
- 14 Moluccan Cockatoo (Cacatua moluccensis), 159
- 15 Pigeon (Columba livia), 179
- 16 Mallard Duck (Anas platyrhynchos), 203
- 17 Red Tailed Hawk (Buteo jamaicensis), 227
- 18 Great Horned Owl (Bubo virginianus), 249

DETAILED CONTENTS

- Radiology Equipment and Positioning Techniques, 1
- 2 Anatomic Art, 17
 Viscera of the Avian Coelom:
 Lateral, 18
 Viscera of the Avian Coelom:
 Ventral, 19
 Avian Skull: Lateral, 20
 Avian Skull: Ventral, 21
 Avian Skeleton: Lateral, 22
 Avian Skeleton: Ventral, 23
- 3 Budgerigar (Melopsittacus undulatus), 25 Radiographic Studies, 26 Viscera of the Coelom: Laterolateral, 26 Viscera of the Coelom: Ventrodorsal, 27
- 4 Peach-Faced Lovebird (Agapornis
 roseicollis), 29
 Radiographic Studies, 30
 Viscera of the Coelom:
 Laterolateral, 30
 Viscera of the Coelom:
 Ventrodorsal, 31
 Radiographic Contrast Studies, 32
 Gastrointestinal Positive Contrast:
 Laterolateral, 32
 Gastrointestinal Positive Contrast:
 Ventrodorsal, 35
- 5 Cockatiel (Nymphicus
 hollandicus), 39
 Radiographic Studies, 40
 Viscera of the Coelom:
 Laterolateral, 40
 Viscera of the Coelom:
 Ventrodorsal, 41
 Radiographic Contrast Studies, 42
 Gastrointestinal Positive Contrast:
 Laterolateral, 00
 Gastrointestinal Positive Contrast:
 Ventrodorsal, 42

- 6 Sun Conure (Aratinga solstitialis), 47
 Radiographic Studies, 48
 Viscera of the Coelom:
 Laterolateral, 48
 Viscera of the Coelom:
 Ventrodorsal, 49
- 7 Senegal Parrot (Poicephalus senegalus), 51 Radiographic Studies, 52 Viscera of the Coelom: Laterolateral, 52 Viscera of the Coelom: Ventrodorsal, 53 Radiographic Contrast Studies, 54 Gastrointestinal Positive Contrast: Laterolateral, 54 Gastrointestinal Positive Contrast: Ventrodorsal, 58 Gastrointestinal Double Contrast: Laterolateral, 62 Gastrointestinal Double Contrast: Ventrodorsal, 64
- 8 African Grey Parrot (Psittacus
 erithacus erithacus), 67
 Radiographic Studies, 68
 Viscera of the Coelom:
 Laterolateral, 68
 Viscera of the Coelom:
 Ventrodorsal, 69
 Skeleton, 70
 Axial Skeleton: Laterolateral, 70
 Axial Skeleton: Ventrodorsal, 71
 Radiographic Contrast Studies, 72
 Positive Contrast Rhinogram:
 Laterolateral, 72
 Positive Contrast Rhinogram:
 Ventrodorsal, 73
- 9 Orange-Winged Amazon Parrot Radiographic Studies (Amazona amazonica), 75 Radiographic Studies, 76 Viscera of the Coelom (Normal Bird): Laterolateral, 76

Viscera of the Coelom (Normal Bird): Ventrodorsal, 77		Head (Post-contrast): Transverse, 123
Viscera of the Coelom (Obese		Ultrasound of Liver, 124
Bird): Laterolateral, 78 Viscera of the Coelom (Obese	10	Blue and Gold Macaw (Ara
Bird): Ventrodorsal, 79		ararauna), 125
Skeleton, 80		Radiographic Studies, 126
Head: Laterolateral, 80		Viscera of the Coelom:
Head: Oblique (30 Degrees), 81		Laterolateral, 126
Head: Ventrodorsal, 82		Viscera of the Coelom:
Axial Skeleton: Laterolateral, 84		Ventrodorsal, 127
Axial Skeleton: Ventrodorsal, 85		Skeleton, 128
Wing: Mediolateral, 86		Head: Laterolateral, 128
Wing: Caudocranial, 87		Head: Ventrodorsal, 129
Pelvic Limb: Mediolateral, 88		Axial Skeleton: Laterolateral, 130
Pelvic Limb: Craniocaudal, 89		Axial Skeleton: Ventrodorsal, 131
Distal Pelvic Limb: Mediolateral, 90		Wing: Mediolateral, 132
Distal Pelvic Limb: Dorsoplantar,		Wing: Caudocranial, 133
91		Pelvic Limb: Mediolateral, 134
Radiographic Contrast Studies, 92		Pelvic Limb: Craniocaudal, 135
Gastrointestinal Positive Contrast:		Radiographic Contrast Studies, 136
Laterolateral, 92		Gastrointestinal Positive Contrast:
Gastrointestinal Positive Contrast:		Laterolateral, 136
Ventrodorsal, 93		Gastrointestinal Positive Contrast:
Gastrointestinal Double Contrast:		Ventrodorsal, 139
Laterolateral, 98		Computed Tomography Study, 142
Gastrointestinal Double Contrast:		Head: Transverse, 142
Ventrodorsal, 100		
Cloacagram (Retrograde, Positive	11	Goffin Cockatoo (Cacatua
and Double Contrast):		goffiniana), 147
Laterolateral, 102		Radiographic Studies, 148
Cloacagram (Retrograde, Positive		Viscera of the Coelom:
and Double Contrast):		Laterolateral, 148
Ventrodorsal, 103		Viscera of the Coelom:
Excretory Urogram: Laterolateral, 106		Ventrodorsal, 149
Excretory Urogram: Ventrodorsal,	12	Sulphur-Crested Cockatoo (Cacatua
108	AT US	galerita), 151
Computed Tomography Studies, 110		Computed Tomography Study, 152
Head: Transverse, 110		Head: Transverse, 152
Coelom: Transverse, 113		110000. 1700000000, 152
Coelom (Post-contrast): Transverse,	13	Umbrella Cockatoo (Cacatua
116		alba), 155
Magnetic Resonance Imaging Studies,		
118		Radiographic Studies, 156
Head: Sagittal, 118		Viscera of the Coelom:
Head (Post-contrast): Sagittal, 120		Laterolateral, 156
Head: Transverse, 121		Viscera of the Coelom:
rieua. iransverse, 121		Ventrodorsal, 157

14	Moluccan Cockatoo (Cacatua moluccensis), 159		Distal Pelvic Limb: Dorsoplantar, 191
	Radiographic Studies, 160		Respiratory Series: Spontaneous
	Viscera of the Coelom (Fasted		Ventilation, 192
	Bird): Laterolateral, 160		Respiratory Series: Manual Assisted
	Viscera of the Coelom (Fasted		Inspiration, 192
	Bird): Ventrodorsal, 161		Respiratory Series: Mechanical
	Viscera of the Coelom (Pellet-fed		Ventilator Assisted Inspiration, 192
	Bird): Laterolateral, 162		Radiographic Contrast Studies, 194
	Viscera of the Coelom (Pellet-fed		Gastrointestinal Positive Contrast:
	Bird): Ventrodorsal, 163		Laterolateral, 194
	Skeleton, 164		Gastrointestinal Positive Contrast:
	Head: Laterolateral, 164		Ventrodorsal:, 197
	Head: Ventrodorsal, 165		Cloacagram (Retrograde, Positive
	Axial Skeleton: Laterolateral, 166		Contrast): Laterolateral, 200
	Axial Skeleton: Ventrodorsal:, 167		Cloacagram (Retrograde, Positive
	Wing: Mediolateral, 168		Contrast): Ventrodorsal, 201
	Wing: Caudocranial, 169		commung. remondrating 201
	Pelvic Limb: Mediolateral, 170	16	Mallard Duck (Anas platyrhynchos)
	Pelvic Limb: Craniocaudal, 171		203
	Distal Pelvic Limb: Mediolateral,		
	172		Radiographic Studies, 204
			Viscera of the Coelom:
	Distal Pelvic Limb: Dorsoplantar,		Laterolateral, 204
	173		Viscera of the Coelom:
	Radiographic Contrast Studies, 174		Ventrodorsal, 205
	Gastrointestinal Positive Contrast:		Skeleton, 206
	Laterolateral, 174		Head: Laterolateral, 206
	Gastrointestinal Positive Contrast:		Head: Ventrodorsal, 207
	Ventrodorsal, 175		Axial Skeleton: Laterolateral:,
	Computed Tomography Study, 176		208
	Head: Transverse, 176		Axial Skeleton: Ventrodorsal:, 209
	THE RESERVE OF THE PARTY OF THE		Wing: Mediolateral, 210
15	Pigeon (Columba livia), 179		Wing: Caudocranial, 211
	Radiographic Studies, 180		Pelvic Limb: Mediolateral, 212
	Viscera of the Coelom:		Pelvic Limb: Craniocaudal, 213
	Laterolateral, 180		Distal Pelvic Limb: Mediolateral,
	Viscera of the Coelom:		214
	Ventrodorsal, 181		Distal Pelvic Limb: Dorsoplantar,
	Skeleton, 182		215
	Head: Laterolateral, 182		Radiographic Contrast Studies, 216
	Head: Ventrodorsal, 183		Gastrointestinal Positive Contrast:
	Axial Skeleton: Laterolateral, 184		Laterolateral, 216
	Axial Skeleton: Ventrodorsal, 185		Gastrointestinal Positive Contrast:
	Wing: Mediolateral, 186		Ventrodorsal, 217
	Wing: Caudocranial, 187		Excretory Urogram (Cutaneous
	Pelvic Limb: Mediolateral, 188		Ulnar Vein Contrast
	Pelvic Limb: Craniocaudal, 189		Administration): Laterolateral,
	Distal Pelvic Limb: Mediolateral, 190		219
	District Line Line, Mediciniera, 150		

Excretory Urogram (Medial Metatarsal Vein Contrast Administration): Laterolateral, 220

Computed Tomography Studies, 221 Head: Transverse, 221 Coelom: Transverse, 225

17 Red Tailed Hawk (Buteo

jamaicensis), 227

Radiographic Studies, 228

Viscera of the Coelom: Laterolateral, 228

Viscera of the Coelom:

Ventrodorsal, 229

Skeleton, 230

Head: Laterolateral, 230

Head: Ventrodorsal, 231

Axial Skeleton: Laterolateral, 232

Axial Skeleton: Ventrodorsal, 233

Wing: Mediolateral, 234

Wing: Caudocranial:, 235

Pelvic Limb: Mediolateral, 235

Pelvic Limb: Craniocaudal, 236

Distal Pelvic Limb: Mediolateral, 237

Distal Pelvic Limb: Dorsoplantar, 238

Radiographic Contrast Studies, 239

Gastrointestinal Positive Contrast:

Laterolateral, 239

Gastrointestinal Positive Contrast: Ventrodorsal, 242 Computed Tomography Study, 245 Coelom: Transverse, 245

18 Great Horned Owl (Bubo

virginianus), 249

Radiographic Studies, 250

Viscera of the Coelom:

Laterolateral, 250

Viscera of the Coelom:

Ventrodorsal, 251

Skeleton, 252

Head: Laterolateral, 252

Head: Ventrodorsal, 253

Axial Skeleton: Laterolateral, 254

Axial Skeleton: Ventrodorsal, 255

Wing: Mediolateral, 256

Wing: Caudocranial, 257

Pelvic Limb: Mediolateral, 258

Pelvic Limb: Craniocaudal, 259

Distal Pelvic Limb: Mediolateral,

260

Distal Pelvic Limb: Dorsoplantar,

261

Radiographic Contrast Studies, 262

Gastrointestinal Positive Contrast:

Laterolateral, 262

Gastrointestinal Positive Contrast:

Ventrodorsal, 265

Computed Tomography Study, 268

Head: Transverse, 268

Radiology Equipment and Positioning Techniques



ANATOMIC REFERENCES

Anatomic drawings demonstrating the spatial relationship of the internal organs are provided in Chapter 2. They should be used as general reference material. Individual organs are not always clearly visualized on all radiographs. There are species variations in the size, shape, and location of internal organs. The radiographic appearance of the viscera is also affected by the birds' reproductive status and digestive tract contents. In the case of this text, the reproductive organs were labeled as "gonads" (versus ovary or testes) if no specific anatomic structure (i.e., the syrinx in male duck) could be identified on the radiograph to indicate the bird's sex.

Images in this text were anatomically labeled to coincide with illustrations from several textbooks, including *Atlas of Avian Anatomy Osteology-Arthrology-Myology* (Chamberlain FW, East Lansing, Mich, 1943, Michigan State College Agricultural Experiment Station), *Atlas of Radiographic Anatomy and Diagnosis of Cage Birds* (Krautwald ME et al., Berlin and Hamburg, 1992, Paul Parey Scientific Publishers), *A Color Atlas of Avian Anatomy* (McLelland J, Philadelphia, 1991, Saunders), *Anatomy of the Domestic Birds* (Nickel R et al., 1977, Berlin and Hamburg, Verlag Paul Parey), and *Atlas of Avian Radiographic Anatomy* (Smith S, Smith B, 1992, Philadelphia, Saunders).

EQUIPMENT FOR RADIOGRAPHIC STUDIES

RADIOGRAPHIC UNITS

It is sometimes erroneously presumed that the relatively small body size of birds allows for the use of low-capacity radiology equipment. However, in order to make high-quality radiographic images of birds, the x-ray generator should be capable of producing at least 300 milliamps (mA), the exposure time capability should be .017 (1/60) second or shorter, the kilovolt peak (kVp) settings should have a range of 40-90 kVp, and kVp settings should be adjustable in 2 kVp increments. Highfrequency x-ray generators are recommended because they produce uniform x-ray output. Radiographic exposure factors are more critical in birds than in mammals. Small variations in x-ray output are very noticeable on avian radiographs, especially those made with lower kVp techniques. Small variations in x-ray tube output are more obvious on avian radiographs than on radiographs of dogs and cats; therefore the x-ray generator must be in excellent condition. Older generators are not recommended because they may have mechanical exposure timers that are less accurate than electronic timers and the x-ray output may vary between exposures with identical settings even though the exposure settings are similar.

Over time, use of x-ray tubes causes pitting of the anode or filament damage, resulting in degradation of x-ray output. Radiographs made with damaged tubes may be diagnostic, but the variation in image quality among studies can affect interpretation, especially in sequential examinations monitoring the response to therapy or the progression of disease (e.g., pneumonia). Lung and air sac image opacities are very susceptible to variations in radiographic techniques. If variables in radiographic techniques are not identified, increased pulmonary or air sac opacity caused by underexposure can be misdiagnosed as pulmonary consolidation or air sac membrane thickening. Overexposure, if unrecognized, may result in a misdiagnosis of a positive response to treatment.

Short exposure times (i.e., .017 [1/60] second or shorter) are essential to minimize motion artifact associated with the rapid respiratory rate and generalized muscular tremors that are common in birds. Short exposure times dictate high mA settings to generate sufficient x-ray output (i.e., milliampseconds [mAs]).

Low kVp techniques (40-60 kVp) are preferred for most film screen systems because they produce a long scale of contrast compared with higher kVp techniques. The ability to make small kVp adjustments is essential to optimize image

detail. Digital radiology systems usually use higher kVp techniques than film screen systems, but the ability to adjust the kVp in 2 kVp increments remains desirable.

The generation of the x-ray beam is initiated by the production of electrons in the x-ray tube filament. Diagnostic radiology tubes (except for dental units) have two filaments (focal spots). Smaller focal spot tubes generally produce superior image detail compared with larger focal spot tubes. Focal spot size and tube x-ray output are inversely related. In order to obtain enhanced detail, the smaller focal spot should be selected. Selection of the smaller focal spot may require manual override of preprogrammed settings. For larger birds, higher mAs techniques may not be possible with the small focal spot.

Tabletop (nongrid) techniques are standard for avian radiography. Grids are typically employed to minimize the deleterious effects of scatter radiation generated by larger patients that have a body thickness greater than 10 cm. Although some birds' bodies are greater than 10 cm, grids are typically not necessary because the air within the air sacs does not generate significant scatter radiation.

The x-ray tube stand should allow for adjustments of the focal film distance (FFD). Adjustment of the FFD is used to make small variations in the effective mAs (x-ray exposure reaching the film). The mAs reaching the film is inversely proportional to the square of the distance from the x-ray source (focal spot). To compensate for the lower mA output of the small focal spot, while maintaining the short exposure time, avian radiographic techniques frequently utilize shorter FFDs than typically used for canine and feline radiography. The alteration of the FFD can also be used to make small adjustments to the effective mAs that cannot otherwise be made. Excessive reduction of the FFD distorts the image; therefore FFDs of less than 30 inches (76 cm) are discouraged.

Many types of equipment have been used for making radiographic images of birds. If patient motion is not a factor (i.e., the patient is anesthetized, thus minimizing the respiratory and muscular motion) and the bird is small (less than 20 gm body weight), dental units may be utilized. However, dental radiography units are generally not well-suited for avian radiography because of their relatively large focal spot size, low mA capacity, lack of adjustable collimation, and inability to make short exposure times. For most birds dental radiography film is too small for imaging the entire head or coelom, and the range of contrast for dental film is much lower than for general radiology film. Nevertheless, some detailed images can be made with dental radiology units and film, especially for examinations of the distal extremities. Portable x-ray radiographic units designed for large animal extremity work are not optimal for the radiographic examination of birds because these units are incapable of producing sufficient mA at the required short exposure times without severely reducing the FFD.

Standard diagnostic radiology equipment in good condition designed for small animal (feline and canine) practice should be utilized for making radiographic images of birds. An Innovet Select X-ray System (Summit Industries, Inc., Chicago) was used to produce the radiographic images in this book.

RADIOGRAPHIC FILM-INTENSIFYING SCREENS

Selection of radiographic film and intensifying screens is based on the speed of the system (i.e., mAs required to produce a high-quality diagnostic image). Film-screen speed and detail are inversely related. Faster systems are usually capable of producing less detail than slower systems. Asymetrix Detail Intensifying Screens (3M Animal Care Products, 3M Center, St. Paul, Minn.) and Ultra Detail Plus or SE + radiographic film (3M Animal Care Products, 3M Center, St. Paul, Minn.) were used to produce the radiographic images in this book. This film-screen combination produced a radiographic system speed of 100 to 350. Table 1-1 summarizes radiographic exposure factors used for creating this text's radiographic images using a tabletop technique. These settings are intended to be

Table • 1-1

Radiographic exposure guidelines for several species of birds.

Body Weight		Film-Screen		
(gm)	Avian Species	Systems*	mAs	kVp
30-100	Budgie, Cockatiel	2	7.5	50
100-300	Senegal, Amazon	2	7.5-9.0	52
100-300	Senegal, Amazon	1	6	44
300-600	Amazon, Cockatoo	1	6	46
600-900	Cockatoo, Macaw	1	6	48
900-1200	Macaw, Red-Tailed Hawk	1	6	50

Focal film distance = 102 cm (40 inches), table top technique *Film-Screen Systems:

- 1. 3M Asymetrix detail green light emitting rare earth screens and 3M Ultra Detail Plus film (3M Animal Care Products, St. Paul MN).
- 2. Asymetrix Detail green light emitting rare earth screen with 3M SE + film (3M Animal Care Products, St. Paul MN).

guidelines and may require modification depending on the x-ray generator, film-screen combination, radiographic film processing, and patient size. Other film-screen combinations of similar speed and resolution can also be used if they are of sufficient detail and speed.

The same technique is usually used for the laterolateral and ventrodorsal coelomic radiographic studies. For radiographic studies of the distal extremities (foot and distal portion of the wing), the kVps used for coelomic radiographic studies are usually reduced by 2 to 3 kVp to prevent overexposure of the distal extremities.

DIGITAL RADIOLOGY SYSTEMS

Digital radiographic image capture (e.g., direct digital, computed radiography) is slowly replacing film-screen systems in veterinary medicine and will eventually predominate. Nonscreen film and high-detail film-screen systems produce images with superior detail compared with digital systems; however, digital systems are capable of producing a higher image contrast range than is possible with film. This results in improved image quality. Other advantages of digital radiography include images that can be electronically manipulated, do not require film processing, and are immediately viewable. In addition, digital systems result in fewer repeat exposures caused by incorrect exposure factors and film processing errors. Digital systems often use higher kVp and mAs techniques (10%-15% higher kVp and mAs) than film-screen systems. Special algorithms are required for avian patients, but they are increasingly available from the manufacturers that produce the digital radiology systems.

THE RADIOGRAPHIC EXAMINATION

PATIENT PREPARATION

Birds utilized in this text were healthy and were fasted before radiographic examinations. Birds weighing less than 100 grams were fasted for 2 hours, and larger birds were fasted for 3 to 5 hours before the radiographic procedure. Avian patients, especially those that are debilitated, are more easily compromised by food deprivation than are mammals. The decision to withhold food in a clinical situation is therefore complex. Even more important than restricting free access to food is the recommendation not to administer nutrients by gavage for 4 hours before the radiographic study, especially in debilitated birds in which crop and proventricular emptying times may be prolonged. The stress associated with manual restraint or anesthesia required for the radiographic examination increases the potential for regurgitation and airway aspiration of the digestive tract contents.

Digestive tract filling affects the radiographic appearance of internal organs. An example of this is included in the section on the Moluccan cockatoo. Postprandial digestive tract distention can displace the liver cranially. A distended proventriculus

often obscures visualization of the spleen. Some of the misdiagnoses associated with failure to recognize the effects of a recent feeding include hepatomegaly and cardiomegaly. Mass lesions, free coelomic fluid, or enlargement of internal organs can also be obscured by ingesta. This is caused by the added opacity of the ingesta, which alters the appearance of the coelomic organs. When two organs of similar opacity are in direct contact, their individual outlines can merge together. The physical characteristics of the ingesta can also variably affect the appearance of the digestive tract. High-fiber (e.g., beans), high-fluid-content (e.g., fruit), and some pelleted diets can produce dramatic digestive tract distention, simulating pathologic conditions. Pelleted diets can sometimes cause the interface between the contrast medium and digestive tract mucosa to be indistinct in digestive tract contrast studies. This can simulate the radiographic appearance of enteritis or excessive intestinal mucus.

Rapid respiratory movements and muscle fasciculations (fine motor movements) are common in birds and can degrade the radiographic detail. The muscle fasciculations can be associated with hypothermia, stress, or a light plane of anesthesia. All of these factors should be addressed before proceeding with the radiographic study.

TIMING THE RADIOGRAPHIC EXPOSURE

The avian respiratory rate is more rapid than that of most mammals, the lungs are nonexpansile, and during both inspiration and expiration air is continuously moving into the pulmonary parenchyma and the air sacs. In general, the effect of the respiratory cycle on the radiographic appearance of the pulmonary parenchyma is less in birds than in mammals because avian lungs are nonexpansile. In some cases, distention of the abdominal air sacs is preferred because the increased air sac distention can improve the visibility of the viscera, especially in birds with large amounts of coelomic fat. Timing the radiographic exposure to coincide with air sac inflation is difficult in nonanesthetized birds as a result of the rapid respiratory rate and inability to directly visualize the respiratory movements through the pelage. If timing the radiographic exposure to coincide with inspiration is not possible, the exposure should coincide with a pause in the respiratory cycle. In the case of intubated anesthetized birds, air sac inflation can be achieved by applying positive pressure to the anesthetic circuit reservoir bag. Positive pressure ventilation of 4 to 6 cm of water is recommended, with 8 to 10 cm of water being the maximum amount for most avian species.

ANESTHESIA

Radiographic studies for which the birds are anesthetized with inhalation gas anesthetics are generally completed in less time and are of higher quality than studies in which the birds are not anesthetized. Anesthetized birds are easily positioned with less physical restraint, and the potential for iatrogenic fractures is minimized. It is also possible to inflate the air sacs

with positive pressure ventilation in the intubated bird. Motion artifacts are also reduced with anesthesia. All birds in this text were healthy, and the majority of the studies were performed using inhalation anesthesia or chemical sedation. Birds seen in clinical practice may be severely debilitated, and general anesthesia may be contraindicated; however, it is recommended whenever the anesthetic procedure is deemed safe.

POSITIONING DEVICES

For production of the images for this text, smaller birds (i.e., less than 100 grams body weight) were positioned directly on the radiographic cassette and secured with masking tape, but an acrylic positioning device (Bird Board, 8205 Alba Ct., Citrus Heights, Calif.) was used to facilitate positioning of larger birds. Many avian positioning devices are commercially available. If positioning devices are interposed between the bird and the film or digital sensor, a small increase in kVp (2-4 kVp) may be required to compensate for x-ray beam filtration caused by the device. The need for exposure compensation is especially necessary with low kVp techniques (40-50 kVp) and thicker positioning devices. Positioning devices placed on the x-ray cassette or used with a digital system increase the object film or sensor distance. The increased object film or sensor distance may decrease image detail, but the magnitude of loss is usually minimal. A modified version of the Bird Board is available that allows for direct contact of the bird and the radiographic cassette (Figure 1-1).

PATIENT POSITIONING

Orthogonal projection radiographs (i.e., two projections made at 90 degrees to each other) are indicated for all radiographic studies unless the patient's condition is compromised and the stress involved with restraint and/or anesthesia is deemed too great to obtain both projections.

The standard avian radiographic study includes laterolateral and ventrodorsal studies of the coelom. For radiographic images of the coelom right lateral and ventrodorsal projections were standard in this text. Pectoral extremity (wing) studies include the mediolateral and caudocranial projections. Given the natural curvature of the skeletal structures of the wing, the laterolateral and ventrodorsal projections of the coelom result in similar-appearing images (mediolateral and lateromedial) of the wing, thus the necessity to make the caudocranial projection of the wing. Manual positioning is required for the caudocranial projection of the wing. The whole body techniques do result in orthogonal projections of the pelvic extremities and therefore do not require additional positioning techniques.

POSITIONING TECHNIQUES FOR LATEROLATERAL AND VENTRODORSAL RADIOGRAPHIC STUDIES OF THE AVIAN HEAD

Radiographic studies of the head include laterolateral and ventrodorsal radiographic projections and, when necessary, oblique views. Small wedges of radiolucent foam may be of assistance for precise positioning of the head. For laterolateral and oblique projections, the patient is placed in a lateral recumbent position. Oblique radiographic projections require rotation of 15 to 30 degrees or less off the straight lateral projection. Oblique projections are described by the point of entrance of the x-ray beam to the point of exit. For ventrodorsal projections, the patient is positioned in dorsal recumbency and tape is applied to the ventral aspect of the rhinotheca so that the maxilla is closer to the cassette. This positioning of the head changes the orientation from a rostrocaudal to a ventrodorsal projection (Figures 1-2 to 1-3).

POSITIONING TECHNIQUE FOR THE LATEROLATERAL RADIOGRAPHIC STUDY OF THE AVIAN COELOM

Before positioning the bird for radiographic examination, strips of paper bandage or masking tape are prepared. Paper tape is less traumatic to the pelage and skin than fabric bandage tape,




Figure 1-1 Figure positioning equipment. A modified version of the traditional "Bird Board" is commercially available and allows for direct contact of the bird and the radiographic cassette.




Figure 1-2 Positioning technique for the laterolateral radiographic study of the avian head. The bird is placed in right lateral recumbency with the head resting on the cassette. Stabilization is achieved using radiolucent tape to secure the maxilla and mandible. Care should be taken to avoid applying tape over the eyelids. The x-ray beam is centered (+) ventral to the eye, and the x-ray beam field (dotted lines) includes the entire head and extends to the cervical region. Use of positioning device is optional. A metallic "R" marker is placed on the radiographic cassette indicating that the right side is dependent.

can be removed easily with minimal residue left on the feathers, and is less radiopaque than fabric tapes. The ends of the tape are folded over on themselves for approximately $\frac{1}{4}$ (0.25) inch to facilitate tape removal when the examination is complete or if patient repositioning is required.

Laterolateral radiographic studies in this text were made with the bird in right lateral recumbency (i.e., the right side of the bird is on the x-ray table or positioning device). The laterolateral positioning technique begins with placing the bird's neck in the cervical restraint portion (guillotine) of the positioning device. The body is then gently moved caudally to reduce curvature of the neck. Further gentle traction is applied to the bird's body by immobilizing the legs with tape or strips of bandage gauze attached to the positioning device. To immobilize the legs, bandage gauze can be looped around the distal tarsometatarsus of each leg and then attached to the cleat on the positioning device. The dependent leg is positioned first, and the contralateral leg is then superimposed on the dependent leg. Occasionally, small pieces of radiolucent sponge are placed between the legs to minimize rotation. Finally, the wings are fully extended dorsally to minimize their superimposition on the cranial dorsal thorax, especially the pulmonary parenchyma. To minimize rotation of the body, the wings should be secured individually using tape. The dependent wing is positioned and taped first. Two pieces of tape are used and crossed at the carpal region of each wing separately. If necessary, another piece of tape can be applied across the wings in the humeral region. The primary flight feathers can be separated to allow increased contact between the tape and the surface of the cassette, positioning device, or table. Rotational malpositioning of the body can be associated with asymmetric leg extension or if excessive pressure is applied to the upper wing when it is secured. A metallic "R" marker is placed adjacent to the bird, indicating that the right side is dependent. Proper positioning is confirmed visually and by palpation of the sternum and vertebral column.

Proper patient positioning is essential because any rotation of the bird's body results in distortion of the coelomic organs and can result in misinterpretation of the radiographic findings. On the laterolateral radiographic studies of the coelom in this text, the extremities are superimposed on each




Figure 1-3 Positioning technique for the ventrodorsal radiographic study of the avian head. The patient is positioned in dorsal recumbency. Radiolucent tape is applied to the ventral aspect of the rhinotheca to hyperextend the maxilla at an angle closer to the radiographic cassette. The x-ray beam (+) is centered between the eyes on the midline, and the x-ray beam field (dotted lines) includes the entire head and extends to the region of the cervical vertebrae. Metallic "R" and "L" markers are placed on the radiographic cassette indicating the laterality of the patient.

other. The authors prefer this method of positioning because it results in less rotation of the bird's body. Because the dependent extremities are closer to the film or digital sensor than the contralateral extremity, the dependent extremities can usually be differentiated. Cortical borders are sharper and the bones are less magnified than in the nondependent extremity. In smaller birds, laterality can be difficult to determine because the magnitude of magnification in the nondependent extremities is minimal. An alternative technique to superimposing the extremities is to position the dependent extremity slightly cranial to the contralateral extremity, but this may result in rotation of the body (Figure 1-4).

POSITIONING TECHNIQUE FOR THE VENTRODORSAL RADIOGRAPHIC STUDY OF THE AVIAN COELOM

Precut strips of paper tape, as described for the laterolateral radiographic study of the coelom, are prepared. For the ventrodorsal projection, the bird is placed on the positioning device in dorsal recumbency. The neck is placed in the guillotine and the head is adjusted to the straight rostrocaudal position. Further gentle traction is applied to the bird's body by extending the pelvic limbs slightly caudally and securing them with gauze to the positioning device's cleats. The wings are fully and symmetrically extended laterally and secured with paper tape. Two pieces of tape are crossed at the carpal region of each wing. The sternum (keel) is palpated to confirm that it is superimposed on the vertebral column. Positions of the extremities (legs and wings) are also evaluated for bilateral symmetry. Metallic "R" and "L" markers should be placed on the radiographic cassette indicating the laterality of the patient (Figure 1-5).

POSITIONING TECHNIQUE FOR THE MEDIOLATERAL RADIOGRAPHIC STUDY OF THE AVIAN THORACIC EXTREMITY (WING)

Positioning the body for a mediolateral radiographic study of the thoracic extremity (wing) is similar to the ventrodorsal coelomic study and results in both wings having a mediolateral

projection. In contrast, the mediolateral view of the wing with the bird in lateral recumbency requires positional modification to reduce superimposition of the wings, and only the nondependent wing results in a true mediolateral view; the dependent wing results in a lateromedial view. When the patient is positioned in dorsal recumbency, the mediolateral radiographic study of the wing is preferred over the lateromedial study of the wing (which would require the patient to be in ventral recumbency) because of decreased object film distance (OFD). Positioning the patient for the mediolateral radiographic study of the wing requires that the patient be placed in dorsal recumbency and the body placed to the edge of the cassette so that the entire wing of interest will be included on the radiographic image (Figure 1-6). If the body of the bird is placed on the center of the radiographic cassette, the film size might be insufficient to include the entire wing. Birds with large wing spans may require that the wing be positioned diagonally across the film cassette to maximize the length of the x-ray field.

POSITIONING TECHNIQUE FOR THE CAUDOCRANIAL RADIOGRAPHIC STUDY OF THE AVIAN THORACIC EXTREMITY (WING)

Lateral or supine (dorsal recumbency) positioning of the patient produces similar radiographic images of the wing (i.e., lateromedial and mediolateral). It is therefore necessary to make a caudocranial image of the wing, which is the orthogonal projection of the mediolateral and lateromedial studies. Positioning the avian patient for the caudocranial projection of the wing presents a unique challenge to the technician or veterinarian. To facilitate patient positioning and decrease the chance of iatrogenic fractures, many avian patients require anesthesia or sedation to make the orthogonal caudocranial projection of the wing. The bird is held in an inverted position with the head directed toward the floor and the long axis of the bird's body perpendicular to the surface of the x-ray table. This results in a caudocranial projection of the wing. The leading edge of the wing is placed on the




Figure 1-4 Positioning technique for the laterolateral radiographic study of the avian coelom. The bird is placed in right lateral recumbency on the positioning device, and the guillotine device is applied to the neck. The bird's body is then gently moved caudally to extend the neck. Further gentle traction is applied to the bird's body by extending the pelvic limbs slightly caudally and securing them with gauze to the positioning device's cleats. The wings are then secured to the positioning device in full extension using radiolucent tape. Rotation of the bird's body can result if excessive pressure is applied to the upper wing when it is secured to the positioning device. All extremities (wings and legs) are positioned so that they are superimposed on their contralateral extremity. This helps to minimize rotation of the bird's body. The x-ray beam (+) is centered on the middle portion of the body, and the x-ray beam field (dotted lines) includes the entire bird for small birds. For medium and large birds, the x-ray field includes the body, proximal extremities, and caudal cervical regions. A metallic "R" marker is placed on the radiographic cassette indicating that the right side is dependent.

cassette and the wing is extended manually while the body or legs are held with the other hand. The appropriate metallic "R" or "L" marker is placed on the radiographic cassette indicating whether the radiographic image is of the right or left wing, respectively (Figure 1-7). A lead apron and gloves are worn for the procedure. The technician's hands should not be in the primary x-ray beam. This caudocranial projection is more difficult to make with direct digital radiology units because it may not be possible to move the x-ray sensor (located under the table top) to the edge of the table. In contrast, the craniocaudal projection of the wing is not practical because the length of the flight feathers precludes close apposition of the bones of the wing with the radiographic cassette. In addition, craniocaudal positioning of the wing results in increased OFD, thereby degrading image quality.

POSITIONING TECHNIQUE FOR THE MEDIOLATERAL RADIOGRAPHIC STUDY OF THE AVIAN PELVIC LIMB (LEG)

Mediolateral radiographic images of the pelvic limbs are made with the patient in right lateral recumbency for the right leg and left lateral recumbency for the left leg. Mediolateral projections are preferred to the lateromedial projections of the pelvic limbs because the OFD is less with the mediolateral projection. Positioning and securing the leg of interest are similar as described for the laterolateral radiographic projection of the coelom, except the contralateral leg is rotated dorsal and pulled caudally to minimize superimposition on the leg of interest. Placement of the bird on the positioning device, application of the guillotine to the neck, and circum-

ferential application of bandage gauze around the distal aspects of tarsometatarsi facilitate application of traction on both legs for positioning purposes. The appropriate metallic "R" or "L" marker is placed on the radiographic cassette indicating whether the radiographic image is of the right or left leg, respectively (Figure 1-8).

POSITIONING TECHNIQUE FOR THE CRANIOCAUDAL RADIOGRAPHIC STUDY OF THE AVIAN PELVIC LIMB (LEG)

Craniocaudal radiographic images of the pelvic limbs are made with the patient in a supine (dorsal recumbency) position. Craniocaudal projections of the pelvic limb are optimal as a result of the reduction in OFD. Positioning and securing the leg of interest are similar as described for ventrodorsal radiographic projection of the coelom. Placement of the bird on the positioning device, application of the guillotine to the neck, and circumferential application of bandage gauze around the distal aspect of the tarsometatarsus facilitate application of traction on the leg of interest. The appropriate metallic "R" or "L" marker is placed on the radiographic cassette indicating whether the radiographic image is of the right or left leg, respectively (Figure 1-9).

POSITIONING TECHNIQUES FOR THE MEDIOLATERAL AND DORSOPLANTAR RADIOGRAPHIC STUDIES OF THE AVIAN DISTAL PELVIC EXTREMITY (FOOT)

Mediolateral and dorsoplantar radiographic studies of the foot are made with the patient in the lateral and supine positions, respectively. The mediolateral projection is preferred over the lateromedial study because of decreased OFD. Positioning




Figure 1-5 Positioning technique for the ventrodorsal radiographic study of the avian coelom. The bird is placed on the positioning device in dorsal recumbency. The neck is placed in the guillotine, and gentle traction is applied to the bird's body by extending the pelvic limbs slightly caudally and securing them with gauze to the positioning device's cleats. The wings are secured to the positioning device in full extension using radiolucent tape. Two strips of tape are crossed at the carpal region of each wing. The x-ray beam (+) is centered midline on the caudal portion of the sternum, and the x-ray beam field (dotted lines) includes the coelom, head, and extremities for small birds. For medium and large birds, the x-ray field includes the body, proximal extremities, and caudal cervical regions. Metallic "R" and "L" markers are placed on the radiographic cassette indicating the laterality of the patient.

techniques are similar as described for the pelvic limb radiographic examinations, but the positioning device is not utilized and the foot is secured directly to the film cassette with paper tape. Special attention is paid to separating the digits of the foot to minimize superimposition of the digits. For both projections the toes are spread apart and individual digits secured with individual pieces of radiolucent tape. On the mediolateral view of the foot, superimposition of the phalanges makes it difficult to count them. When interpreting the radiographs, counting the phalanges aids in the identification of the specific digit (i.e., the number of phalange bones in each digit is one greater than the number of the digit itself). Radiopaque numeral markers can be placed adjacent to the digits to help identify them on the radiographs. The markers are taped to the surface of the film cassette with radiolucent tape (Figures 1-10 and 1-11).

RADIOGRAPHIC CONTRAST STUDIES OF THE GASTROINTESTINAL TRACT

The rapid digestive tract transit time of birds and the anatomy of their digestive tract make it possible to study the majority of the digestive tract using a single radiographic protocol. The digestive tract from the esophagus to the large intestine can be radiographically evaluated by administering contrast medium orally. The term "UGI" (upper gastrointestinal) tract contrast study is a misnomer in birds because the colon, and occasionally the cloaca, is also evaluated on these studies. Evaluation of the cloaca is best performed with retrograde studies, wherein the contrast medium is administered directly into the cloaca.

Ideally, the ingluvies (crop) and proventriculus should be empty before beginning a digestive tract contrast medium study, although the effects of food deprivation must be considered. A fasting period before administration of the contrast medium is recommended, and the duration of the fasting period depends on the metabolic requirements of the bird and its overall health. Smaller birds do not tolerate food restriction as well as larger birds because of their higher metabolic rate. Recommendations for food restriction are provided in the discussion on patient preparation. The presence of food in the ingluvies (crop) decreases the volume of contrast medium that can be safely administered. Ingesta in the gastrointestinal tract also degrades the detail of the interface between contrast medium and the digestive tract mucosa by absorbing the contrast medium and impeding full contact of the contrast medium with the digestive tract mucosa. Ingesta may also delay passage




Figure 1-6 Positioning technique for the mediolateral radiographic study of the avian thoracic extremity (wing) with the patient in dorsal recumbency. Positioning of the body for the mediolateral radiographic study of the wing is similar to that described for the ventrodorsal coelomic study, except the body is positioned to the side of the cassette so that the entire wing of interest will be included on the radiographic image. The wing is fully extended laterally from the pectoral girdle and taped directly to the radiographic cassette. Immobilization of the wing is best achieved by crossing the tape in the region of the carpus. Additional tape can be applied to the proximal and distal aspects of the wing if necessary. The x-ray beam (+) is centered in the mid-diaphyseal region of the radius and ulna. The x-ray beam field (dotted lines) encompasses the entire wing, including the scapulo-humeral joint. Metallic "R" and "L" markers are placed on the radiographic cassette indicating the laterality of the patient.


of the contrast medium through the digestive tract. Hyperalimentation preparations are particularly unpredictable in the way they mix with contrast media. Pelleted foods also affect the appearance of the intestinal mucosal contrast medium interface. Birds eating pelleted diets often have an indistinct intestinal contrast medium mucosal interface on radiographs. The cause of this is unknown. It is recommended that birds weighing more than 300 grams not be fed pelleted diets within 4 hours of digestive tract contrast studies if their health status allows such a fasting period. This general recommendation is based on clinical experience, and currently no scientific studies have been performed to validate this theory.

Survey ventrodorsal and right laterolateral radiographs are made immediately before digestive tract contrast studies are initiated. The caudal cervical area and the entire coelom are included on the images. These survey radiographs provide information regarding the distention and location of the digestive tract and are essential for confirming that the radio-

graphic exposure factors are optimized. Radiographs made on previous days are not sufficient to serve as survey radiographs because they may not reflect the current status of the digestive tract. As a general rule, positive contrast medium studies utilize slightly higher kVp settings (2-4 kVp) than those kVp settings used for survey radiographs. This compensates for the increased opacity of the contrast medium.

The contrast medium used for the digestive tract studies in this text was barium sulfate 30% weight to volume (w/v). The contrast medium can be warmed to a temperature slightly greater than room temperature by immersing the syringe containing the contrast medium in warm water. Just prior to contrast medium administration, the liquid in the syringe should be mixed and tested to check the temperature. Warming the contrast medium decreases abnormal peristalsis and hypothermia caused by chilled contrast medium. The dose of contrast medium/gram body weight varies among species of birds. A very broad recommendation would be

Figure 1-7 Positioning technique for the caudocranial radiographic study of the avian thoracic extremity (wing). The anesthetized bird is held in an inverted position with the head directed toward the floor and long axis of the bird's body perpendicular to the surface of the x-ray table. The wing to be imaged is fully extended, and the cranial (leading) edge of the wing is placed on the film cassette. The bird is angled so that the x-ray beam is aligned in a true caudocranial direction through the wing. The x-ray beam (+) is centered in the middle of the wing, which is usually the mid-diaphyseal region of the radius and ulna. The x-ray beam field (dotted lines) encompasses the entire wing, including the scapulo-humeral joint. Care should be exercised so that the technician's hands are outside of primary x-ray beam. The appropriate metallic "R" or "L" marker is placed on the radiographic cassette indicating whether the radiographic image is of the right or left wing, respectively.



25 to 50 ml/kg body weight. An alternative method of dose calculation is to estimate the volume of food that could be safely administered to the patient via crop gavage and administer 50% to 75% of this amount of contrast medium. Species variation in crop and proventricular volume and the presence of generalized disease or digestive tract pathology, including suspected obstruction, necessitate reducing the amount of contrast medium administered. The survey radiographs made just prior to the administration of contrast medium provide information regarding the digestive tract volume and mitigating circumstances that could affect the volume of contrast medium administered. Hygroscopic water-soluble contrast media (e.g., diatrizoate meglumine and diatrizoate sodium, Gastrografin®, Bracco Pharmaceuticals, Princeton, N.J.) used in mammals are not recommended for radiographic studies of the digestive tract of birds because they can precipitate severe metabolic and fluid imbalances. In addition, the mucosal detail using hygroscopic water-soluble contrast media is less than with barium sulfate. The double contrast studies utilize a combination of positive contrast medium and air. These studies are intended to speed the progression of the contrast medium through the digestive tract. Double contrast studies may also produce superior proventricular mucosal detail. It is important for the clinician to be aware of the potential for regurgitation of the contrast material that may occur as a result of the increased volume of room air in the gastrointestinal tract.

For the radiographic studies in this text, the contrast medium was administered via a rigid or soft gavage tube passed into the crop. Before administering the contrast medium, the position of the gavage tube was verified by palpation to ensure that the tube was not intratracheal. The right cervical area can be palpated as

the tube is passed. If the tube is within the trachea, the gavage tube will not be palpable as a separate entity from the trachea. While the contrast material is being administered, the bird's oral cavity should be visually monitored constantly to ensure that fluid is not accumulating in the oropharynx.

For the digestive tract contrast studies published in this text, birds were anesthetized to facilitate administration of the contrast medium, optimize patient positioning, and reduce handling. These concepts also apply to clinical patients; however, the overall health of the patients should be evaluated to determine if anesthesia is safe. In general, it is recommended that birds be anesthetized for the survey radiographs, then administered the contrast medium while still under anesthesia and subsequent images acquired before the patient is allowed to recover from anesthesia. In our practice we typically acquire the survey, 0.25 and 0.5 hour radiographic images under anesthesia if the patient's health allows. Tracheal intubation to maintain anesthesia should be utilized to minimize aspiration of contrast material while the series of radiographic images are acquired for the study. Anesthesia of smaller patients (e.g., lovebirds, cockatiels) can be maintained utilizing an anesthetic cone. Patients of this size are typically not intubated because of the risk of tracheal mucosal irritation, and subsequent formation of a transluminal membrane which outweighs the risk of contrast medium aspiration. For both the intubated and the nonintubated birds, the cranial portion of the bird's body can be elevated by securing the bird to the acrylic positioning device and elevating the cranial end of the board by sliding an object such as a sandbag under it. Elevation of the cranial portion of the bird's body can minimize retrograde flow of the contrast medium into the oral cavity. Minimizing retrograde flow of contrast medium into the oral cavity can also be




Figure 1-8 Positioning technique for the mediolateral radiographic study of the avian pelvic limb. The bird is positioned in lateral recumbency on the positioning device so that the limb of interest is dependent and fully extended. Radiolucent tape is applied to the distal tarsometatarsus and digits of the leg of interest. The contralateral limb is pulled caudally and dorsally to minimize superimposition. In order to apply enough traction to the contralateral limb, gauze bandage is wrapped around the distal tarsometatarsus and secured with tape. The x-ray beam is centered (+) on the mid-diaphyseal region of the tibiotarsal bone, and the x-ray beam field (dotted lines) includes the entire limb of interest (including the coxofemoral joint). The appropriate metallic "R" or "L" marker is placed on the radiographic cassette indicating whether the radiographic image is of the right or left leg, respectively.

achieved by placing loose elastic bandage material around the bird's neck to partially occlude the cervical esophagus, but care should be exercised to ensure that the bandage does not compress the trachea. During the entire procedure and immediately before extubation, the oral cavity should be monitored for the presence of regurgitated contrast medium. The impact of anesthetic protocols on gastrointestinal transit time has not been scientifically studied in avian patients; however, given their rapid digestive tract transit time, the impacts appear to be minimal in healthy avian patients. It is important to note that in some birds, hypersensitivity to the gas anesthetic may

induce vomiting and the bird may vomit upon recovery from the anesthetic event.

Contrast medium regurgitation and subsequent tracheal aspiration can occur when performing digestive tract contrast studies and is most common when larger volumes of contrast medium are administered. The authors find regurgitation to be less frequent in anesthetized birds than non-anesthetized birds. If contrast medium regurgitation does occur and the bird is anesthetized, attempts to remove the contrast medium from the oropharynx should be immediately initiated to minimize tracheal aspiration and reflux of




Figure 1-9 Positioning technique for the craniocaudal radiographic study of the avian pelvic limb. The bird is positioned in dorsal recumbency on the positioning device. Gauze bandage is wrapped around the distal tarsometatarsus and traction applied to fully extend the pelvic limb. The toes are individually secured using radiolucent tape. The x-ray beam is centered (+) on the mid-diaphyseal region of the tibiotarsal bone, and the x-ray beam field (dotted lines) includes the entire limb of interest (including the coxofemoral joint). The appropriate metallic "R" or "L" marker is placed on the radiographic cassette indicating whether the radiographic image is of the right or left leg, respectively.

contrast medium through the choanal slit into the nasal cavity. Cotton-tipped applicators can be used to remove the contrast medium from the mouth. Contrast medium aspirated into the trachea can be identified on radiographs. In addition to seeing contrast material in the trachea, it may also be visualized in the major bronchi and air sacs, most commonly the left caudal thoracic air sac. Small volumes of aspirated contrast medium are usually well tolerated; however, larger volumes may result in bronchial obstruction. Contrast medium in the pulmonary parenchyma is rarely identified and seen only in cases in which the bird has aspirated a large volume of material. If the contrast medium is aspirated into the lungs, it is usually associated with severe respiratory distress.

Digestive tract transit time is generally more rapid in birds than mammals, especially in the smaller birds and the frugivorous species. In order to perform a complete digestive tract study, each segment of the digestive tract (i.e., cervical esophagus, crop, thoracic esophagus, proventriculus, ventriculus, small intestines, large intestines, and occasionally the cloaca when it is distended with contrast medium) should be documented. Documentation of proventricular emptying is also helpful.

Timing of the radiographic exposures for digestive tract radiographic studies is based on many factors, but all studies

should include laterolateral and ventrodorsal projections made immediately after the administration of the contrast medium, especially if the esophagus, crop, and proventriculus are of primary interest. In some species (i.e., ducks and geese), the gastrointestinal transit times are extremely rapid and the contrast medium may have already entered the small intestines on the first set of radiographs. In such a case, additional radiographs are then made at 15- to 30-minute intervals and then hourly. Because of the variation in digestive tract transit times of different species of birds, the contrast material, and the effects of systemic disease, it is very difficult to propose rigid guidelines for timing the exposures. The timing is usually determined by evaluating the progress of the contrast medium on the initial set of radiographs. If contrast medium has entered the intestines on the immediate postcontrast administration radiographs, the next series of radiographs are taken in 15 minutes. If only a small volume of contrast medium has exited the crop, the next series of radiographs are taken 30 to 60 minutes postcontrast medium administration. Ventrodorsal and laterolateral radiographs are then made every 30 minutes until contrast medium has entered the colon.

Double contrast digestive tract studies usually have shorter transit times and produce superior mucosal detail than is seen on the positive contrast study. Because the crop and proventriculus are more fully distended on double contrast proce-




Figure 1-10 Positioning technique for the mediolateral radiographic study of the avian distal pelvic extremity (foot). The bird is placed in lateral recumbency with the leg of interest in the dependent position. The distal portion of the leg is taped at the distal aspect of the tarsometatarsus. The digits of the foot are also secured with individual pieces of radiolucent tape. Radiopaque numeral markers can be placed adjacent to the digits to assist in their identification on the radiograph. Radiolucent tape is preferred over gauze because it allows more precise positioning of the digits. The nondependent foot is extended caudally to separate the feet and minimize superimposition. The kVP is slightly reduced (i.e., 2-4 kVP) from the technique used for the laterolateral radiographic examination of the coelom, to prevent overexposure of the digits. The x-ray beam is centered (+) on the condyles of the tarsometatarsal bone, and the x-ray beam field (dotted lines) includes all of the phalanges. The appropriate metallic "R" or "L" marker is placed on the radiographic cassette indicating whether the radiographic image is of the right or left foot, respectively.

dures, the double contrast procedures may be better suited for evaluation of mural thickness and mucosal patterns of these organs than the single contrast studies. Double contrast procedures usually require anesthesia because the gas infused into the crop will be regurgitated in most awake birds. If air is immediately expelled, it is acceptable to administer additional volumes of air to distend the crop. Because of the smaller amount of contrast medium used and the faster time in which the contrast medium evacuates the crop, the potential for contrast medium aspiration into the respiratory tract may be less than in the positive contrast studies.

Optimal evaluation of hollow organs requires that they be fully distended at the time of image capture. Orally administered contrast medium does not predictably result in cloacal distention. Therefore retrograde administration of contrast medium (i.e., via the vent) is required. Before administration

of the contrast medium, the cloaca can be gently flushed with isotonic saline. Positive and double contrast procedures can be performed. The double contrast cloacagram usually follows the positive contrast cloacagram and requires removal of the positive contrast medium before administration of room air or carbon dioxide. Carbon dioxide is favored over room air to minimize the potential for intravascular air embolization. Removal of positive contrast medium pooled in the cloaca should be performed before the negative contrast (i.e., room air or carbon dioxide) is introduced. This facilitates visualization of the cloacal mucosal surface. Theoretically, it is possible to reflux fecal matter into the ureters when performing retrograde cloacograms, but this phenomenon has not been recognized in our studies.

Because radiographic images are two-dimensional representations of three-dimensional objects, additional studies




Figure 1-11 Positioning technique for the dorsoplantar radiographic study of the avian distal pelvic extremity (foot). The bird is placed in dorsal recumbency, and the distal portion of the leg is secured with radiolucent tape. The digits of the foot are fully extended and separated using individual pieces of radiolucent tape. The kVP is reduced slightly (i.e., 2-4 kVP) from the technique used for the ventrodorsal radiographic examination of the coelom to prevent overexposure of the digits. The x-ray beam is centered (+) on the condyles of the tarsometatarsal bone, and the x-ray beam field (dotted lines) includes all of the phalanges. The appropriate metallic "R" or "L" marker is placed on the radiographic cassette indicating whether the radiographic image is of the right or left foot, respectively.

may be required to fully assess the motility of the digestive tract. Flouroscopy can be used to evaluate proventricular and ventricular motility, but this requires specialized equipment.

RADIOGRAPHIC CONTRAST STUDIES OF THE URINARY TRACT

Survey radiographs are essential for establishing the radiographic technique and identifying subtle radiographic changes in the avian urinary tract after administration of contrast. Radiographic exposures should be made immediately after injection of the contrast medium. The contrast medium should be injected as a bolus directly into the intravenous catheter. Our clinical experience has led us to use nonionic iodinated contrast media for excretory urography. We have not performed double blind studies, but the nonionic agents seem to be associated with fewer anesthetic issues and less nausea and regurgitation. A general recommendation for the volume of contrast medium to administer is to calculate 50% of the recommended mammalian dose/kg body weight. Extremely concentrated products are not recommended because of their potential hyperosmotic effects. The selection

of contrast medium is also affected by the volume of contrast medium required.

Because of the renal portal system, the injection sight dramatically affects urinary tract opacification. Injection of the contrast material into the medial metatarsal vein results in superior ureteral opacification compared with cutaneous ulnar vein or jugular vein injections. The contrast medium delivered to the renal parenchyma by the renal portal system after being injected into the medial metatarsal vein is less dilute compared with when it is injected into other cranially located peripheral veins. Thus administration of contrast material via the medial metatarsal vein produces a superior urogram, but the authors have not performed studies to document the safety of this route of administration of contrast medium.

COMPUTERIZED TOMOGRAPHY (CT) AND MAGNETIC RESONANCE IMAGING (MRI)

The advantages of advanced imaging modalities, commonly accepted in mammalian imaging, are less obvious in avian patients. This is primarily due to the small body size of birds. Resolution depends greatly on the quality of the equipment

and software. All of the advanced imaging modalities require general anesthesia because image acquisition time can be prolonged.

Computed tomography (CT) and magnetic resonance imaging (MRI) are very helpful for identifying the origin of mass lesions and complex skull abnormalities, but are limited in their capability to evaluate subtle soft tissue and skeletal lesions in birds. CT is very effective for identifying focal pulmonary lesions if the size of the patient and the size of the lesion are sufficient for detection. MRI has been very useful in evaluating the diverticuli of the infraorbital sinus. Threedimensional CT reconstructions are especially helpful in evaluating the sinuses and skull bones.

Birds potentially undergoing MRI examinations should be radiographed to verify that they do not have internal foreign

metallic material within their digestive tract or elsewhere. This is especially true with free-ranging birds. Ferrous metals can migrate through soft tissue planes if placed in the MR field. Contrast medium can be used to enhance information

obtained from CT and MRI studies. In this text a lower dose of contrast medium than is recommended for mammals was used for the CT studies. For MRI studies the contrast medium should be injected directly into the intravenous catheter and not diluted. This practice delivers a more concentrated bolus of contrast medium to the patient than would occur if the contrast medium were injected into the intravenous extension tubing. Further scientific evaluation of optimal doses and safety is required.

Anatomic Art






Figure 2-1 Anatomic drawing (view of the left side) of the viscera of an adult bird.

- Trachea
 Syrinx
 Aorta

- 4. Lung
 5. Pulmonary vein
 6. Proventriculus
 7. Gonad

- 8. Spleen
- 9. Kidney
- 10. Colon
- 11. Cloaca

- 12. Cervical esophagus13. Crop14. Thoracic esophagus15. Brachiocephalic artery and aorta16. Polymerus artery
- 16. Pulmonary artery
 17. Heart
 18. Liver

- 19. Proventricular-ventricular isthmus
- 20. Ventriculus
- 21. Intestines




Figure 2-2 Anatomic drawing (ventrodorsal view) of the viscera of an adult bird.

- 1. Trachea
 2. Syrinx
 3. Heart base vessel
 4. Heart
 5. Proventriculus

- 6. Liver 7. Ventriculus 8. Colon
- 9. Cervical esophagus

- 10. Crop
 11. Thoracic esophagus
 12. Brachiocephalic artery and aorta
 13. Lung
 14. Kidneys
 15. Pancreas
 16. Duodenum
 17. Intestines

- 17. Intestines 18. Cloaca




Figure 2-3 Anatomic drawing (view of the left side) of the skull of an adult bird.

- Keratinized maxillary beak
 Premaxillary bone
- 3. External nare
- 4. Craniofacial flexion zone
- 5. Frontal bone
 6. Orbit
 7. Cranium

- 8. Jugal [zygomatic] bone

- 9. Parietal bone
- 10. Quadrate bone
- 11. Keratinized mandibular beak
- 12. Palatine bone

- 12. Falathle bolle13. Mandible14. Hyoid bones15. Pterygoid bone




Figure 2-4 Anatomic drawing (ventral view) of the skull of an adult bird.

- Premaxillary bone
 Scleral ossicles
 Hyoid bones
 Jugal (zygomatic) bone
 Quadrate bone
 Keratinized maxillary beak

- 7. Orbit8. Mandible9. Pterygoid bone10. Cranium11. Palatine bone




Figure 2-5 Anatomic drawing (view of the left side) of the skeleton of an adult bird.

- 1. Cervical vertebra
- 2. Humerus
- 3. Thoracic vertebra
- 4. Scapula
- 5. Rib
- 6. Uncinate process of rib 7. Ilium
- 8. Synsacrum
- 9. Head of femur
- 10. Femur
- 11. Obturator foramen
- 12. Ilioischiadic foramen

- 13. Ischium14. Caudal vertebra
- 15. Pygostyle
- 16. Clavicle 17. Coracoid
- 18. Sternum
- 19. Costochondral junction of rib 20. Patella
- 21. Fibula
- 22. Pubic bones23. Tibiotarsal bone





Figure 2-6 Anatomic drawing (ventral view) of the skeleton of an adult bird.

- 1. Cervical vertebra
- 2. Coracoid
- 3. Head of humerus
- 4. Ventral tubercle of humerus
 5. Scapula
- 6. Thoracic vertebra
- 7. Head of femur 8. Synsacrum
- 9. Caudal vertebrae 10. Clavicle
- 11. Dorsal tubercle of humerus

- 12. Sternum
- 13. Rib 14. Ilium
- 15. Trochanter of femur
 16. Femur
- 17. Fibula
- 18. Ischium
- 19. Pubic bone 20. Tibiotarsal bone 21. Pygostyle

Budgerigar (Melopsittacus undulatus)






Figure 3-1, A-B

Type of Bird: Budgerigar Type of Study: Viscera of the coelom

Projection: Laterolateral (right lateral

recumbency) Weight of Bird: 30 g Gender: Unknown Reproductive Status: Intact

Age: Adult

- 1. Trachea
- 2. Crop
- 3. Brachiocephalic artery and aorta
- 4. (Brachiocephalic artery)
- 5. (Aorta)
 6. Pulmonary artery
- 7. Pulmonary vein
- 8. Heart
- 9. Left atrium
- 10. (Esophagus)
- 11. Liver
- 12. Lung
- 13. Syrinx
- 14. (Gonad) 15. (Ovary) 16. (Testes)

- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. Cloaca
- 21. (Cervical air sac)
- 22. (Clavicular air sac)
- 23. Thoracic air sac 24. (Abdominal air sac)
- 25. Apex of heart
- 26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys
- 28. (Spleen)

NOTE: Structures in parentheses are not





Figure 3-2, A-BType of Bird: Budgerigar
Type of Study: Viscera of the coelom Projection: Ventrodorsal Weight of Bird: 30 g Gender: Unknown Reproductive Status: Intact Age: Adult

- 1. Trachea

- Cervical air sac)
 Crop
 Clavicular air sac
- 5. Brachiocephalic artery and aorta6. Heart base vessel
- 7. Heart 8. Lung
- 9. Thoracic air sac
- 10. Liver

- 11. Kidneys12. Proventriculus13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. (Cloaca)

Peach-Faced Lovebird (Agapornis roseicollis)






Figure 4-1, A-B

Type of Bird: Peach-Faced Lovebird Type of Study: Viscera of the coelom Projection: Laterolateral (right lateral recumbency)

Weight of Bird: 50 g

Gender: Unknown Reproductive Status: Intact

Age: Adult

- 1. Trachea
- 2. (Crop)3. Brachiocephalic artery and aorta
- 4. (Brachiocephalic artery)
- 5. (Aorta)
- 6. Pulmonary artery
- 7. Pulmonary vein 8. Heart
- 9. Left atrium
- 10. Esophagus
- 11. Liver
- 12. Lung
- 13. Syrinx
- 14. Gonad
- 15. (Ovary) 16. (Testes)

- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. (Cloaca)
- 21. (Cervical air sac)
 22. (Clavicular air sac)
- 23. Thoracic air sac
- 24. Abdominal air sac
- 25. Apex of heart
- 26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys
- 28. (Spleen)




Figure 4-2, A-BType of Bird: Peach-Faced Lovebird Type of Study: Viscera of the coelom Projection: Ventrodorsal Weight of Bird: 50 g Gender: Unknown Reproductive Status: Intact

Age: Adult

- 1. Trachea
- 2. (Cervical air sac)

- (Crop)
 Clavicular air sac
 (Brachiocephalic artery and aorta)
- 6. (Heart base vessel)
- 7. Heart
- 8. Lung
 9. Thoracic air sac
- 10. Liver

- 11. Kidneys12. (Proventriculus)
- 13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. (Cloaca)




Figure 4-3, A-B

Type of Bird: Peach-Faced Lovebird
Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)
3 ml administered via gavage tube
Projection: Laterolateral (right lateral

recumbency) Weight of Bird: 50 g Gender: Unknown
Reproductive Status: Intact

Age: Adult

lmage	Time (hr)
A	Scout
В	0.25

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 4-3, C-DType of Bird: Peach-Faced Lovebird
Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 3 ml administered via gavage tube Projection: Laterolateral (right lateral

recumbency)
Weight of Bird: 50 g
Gender: Unknown Reproductive Status: Intact Age: Adult

lmage	Time (hr)
С	2.0
D	4.0

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 4-3, E

Type of Bird: Peach-Faced Lovebird
Type of Study: Gastrointestinal positive
contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 3 ml administered via gavage tube Projection: Laterolateral (right lateral recumbency)

Weight of Bird: 50 g Gender: Unknown Reproductive Status: Intact

Age: Adult

Image	Time (hr)
Е	7.5

- 1. Crop
- 2. (Esophagus)
- 3. Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. Large intestine
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 4-4, A-BType of Bird: Peach-Faced Lovebird
Type of Study: Gastrointestinal positive contrast study

Contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 3 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 50 g Gender: Unknown Reproductive Status: Intact Age: Adult

Image	Time (hr)
A	Scout
В	0.25

- 1. Crop
- 2. Esophagus3. Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)
 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 4-4, C-D

Type of Bird: Peach-Faced Lovebird Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 3 ml administered via gavage tube Projection: Ventrodorsal Weight of Bird: 50 g

Gender: Unknown Reproductive Status: Intact Age: Adult

lmage	Time (hr)
С	2.0
D	4.0

- Crop
 Esophagus
 Proventriculus
 Ventriculus
- $5. \ Intestines$
- 6. Duodenum
- 7. (Large intestine)
- 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 4-4, E

Type of Bird: Peach-Faced Lovebird Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 3 ml administered via gavage tube Projection: Ventrodorsal Weight of Bird: 50 g


Gender: Unknown Reproductive Status: Intact

Age: Adult

Image	Time (hr)
Е	7.5

- (Crop)
 (Esophagus)
- 3. Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. Large intestine
- 8. (Cloaca)
- 9. Proventricular-ventricular isthmus

Cockatiel (Nymphicus hollandicus)






Figure 5-1, A-B

Type of Bird: Cockatiel Type of Study: Viscera of the coelom Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 98 g

Gender: Male

Reproductive Status: Intact

Age: Adult

- 1. Trachea
- Crop
 Brachiocephalic artery and aorta
- 4. (Brachiocephalic artery)
- 5. Àorta
- 6. Pulmonary artery
- 7. Pulmonary vein
- 8. Heart
- 9. Left atrium
- 10. Esophagus
- 11. Liver
- 12. Lung
- 13. Syrinx
- 14. Gonad
- 15. (Ovary) 16. (Testes)

- $17.\ Proventriculus$
- 18. Ventriculus
- 19. Intestines
- 20. Cloaca
- 21. (Cervical air sac)
 22. (Clavicular air sac)
- 23. Thoracic air sac 24. Abdominal air sac
- 25. Apex of heart
- 26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys
- 28. Spleen




Figure 5-2, A-B
Type of Bird: Cockatiel
Type of Study: Viscera of the coelom
Projection: Ventrodorsal
Weight of Bird: 98 g
Gender: Male
Reproductive Status: Intact Reproductive Status: Intact Age: Adult

- Trachea
 Cervical air sac
 Crop
- 4. Clavicular air sac
- 5. Brachiocephalic artery and aorta6. Heart base vessel7. Heart

- 8. Lung
 9. Thoracic air sac
- 10. Liver

- 11. Kidneys12. (Proventriculus)
- 13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. (Cloaca)




Figure 5-3, A-BType of Bird: Cockatiel
Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 5 ml administered via gavage tube Projection: Ventrodorsal Weight of Bird: 98 g

Gender: Male Reproductive Status: Intact Age: Adult

lmage	Time (hr)
A	Scout
В	0.25

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
 7. (Large intestine)
- 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 5-3, C-D
Type of Bird: Cockatiel

Type of Study: Gastrointestinal positive contrast study
Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)

5 ml administered via gavage tube Projection: Ventrodorsal Weight of Animal: 98 g Gender: Male Reproductive Status: Intact Age: Adult

lmage	Time (hr)
С	0.5
D	2.0

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
 7. Large intestine
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 5-4, A-BType of Bird: Cockatiel

Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 5 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 98 g Gender: Male Reproductive Status: Intact Age: Adult

Image	Time (hr)
A	Scout
В	0.25

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)
- 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 5-4, C-DType of Bird: Cockatiel

Type of Study: Gastrointestinal positive contrast study


Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)
5 ml administered via gavage tube
Projection: Ventrodorsal
Weight of Bird: 98 g
Gender: Male


Reproductive Status: Intact Age: Adult

lmage	Time (hr)
С	0.5
D	2.0

- Crop
 Esophagus
- 3. Proventriculus 4. Ventriculus
- 5. Intestines6. Duodenum
- 7. Large intestine
- 8. Cloaca
- 9. Proventricular-ventricular isthmus

Sun Conure (Aratinga solstitialis)








Figure 6-1, A-B
Type of Bird: Sun Conure Type of Study: Viscera of the coelom Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 126 g
Gender: Unknown

Reproductive Status: Intact

Age: Adult

- 1. Trachea
- 2. Crop
- 3. Brachiocephalic artery and aorta
- 4. Brachiocephalic artery
- 5. Aorta
- 6. Pulmonary artery
- 7. Pulmonary vein
- 8. Heart
- 9. Left atrium
- 10. Esophagus
- 11. Liver
- 12. Lung
- 13. Syrinx
- 14. (Gonad) 15. (Ovary) 16. (Testes)

- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. (Cloaca)
- 21. (Cervical air sac)
 22. (Clavicular air sac)
- 23. Thoracic air sac
- 24. Abdominal air sac
- 25. (Apex of heart)26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys
- 28. Spleen




Figure 6-2, A-B Type of Bird: Sun Conure
Type of Study: Viscera of the coelom Projection: Ventrodorsal Weight of Bird: 126 g Gender: Unknown Reproductive Status: Intact Age: Adult


- 1. Trachea

- (Cervical air sac)
 Crop
 Clavicular air sac
 Brachiocephalic artery and aorta
- 6. Heart base vessel 7. Heart

- 8. Lung
 9. Thoracic air sac
 10. Liver

- 11. Kidneys12. (Proventriculus)13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. Cloaca

Senegal Parrot (Poicephalus senegalus)






Figure 7-1, A-B

Type of Bird: Senegal Parrot Type of Study: Viscera of the coelom Projection: Laterolateral (right lateral

recumbency) Weight of Animal: 146 g

Gender: Unknown

Reproductive Status: Intact

Age: Adult

- 1. Trachea
- 2. Crop
- 3. (Brachiocephalic artery and aorta)4. Brachiocephalic artery
- 5. Aorta
- 6. Pulmonary artery
- 7. Pulmonary vein
- 8. Heart
- 9. Left atrium
- 10. Esophagus
- 11. Liver
- 12. Lung
- 13. Syrinx
- 14. Gonad 15. (Ovary)
- 16. (Testes)

- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. Cloaca
- 21. (Cervical air sac)
- 22. (Clavicular air sac)
- 23. Thoracic air sac
- 24. Abdominal air sac 25. Apex of heart
- 26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys
- 28. Spleen




Figure 7-2, A-B
Type of Bird: Senegal Parrot
Type of Study: Viscera of the coelom
Projection: Ventrodorsal
Weight of Animal: 146 g Gender: Unknown Reproductive Status: Intact Age: Adult

- Trachea
 (Cervical air sac)
 (Crop)
 Clavicular air sac
 (Brachiocephalic artery and aorta)
- 6. (Heart base vessel)
- 7. Heart
- 8. Lung
 9. Thoracic air sac

- 10. Liver11. Kidneys12. (Proventriculus)
- 13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. Cloaca




Figure 7-3, A-B

Type of Bird: Senegal Parrot
Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 7 ml administered via gavage tube

Projection: Laterolateral (right lateral recumbency)

Weight of Animal: 146 g Gender: Unknown Reproductive Status: Intact Age: Adult

lmage	Time (hr)
A	Scout
В	0.5

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
- 8. Cloaca
- 9. Proventricular-ventricular isthmus




Figure 7-3, C-D
Type of Bird: Senegal Parrot
Type of Study: Gastrointestinal positive contrast study
Contract Medium Parium gulfate gunner

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 7 ml administered via gavage tube Projection: Laterolateral (right lateral

recumbency) Weight of Animal: 146 g Gender: Unknown

Reproductive Status: Intact Age: Adult

Image	Time (hr)
С	1.5
D	2.5

- Crop
 Esophagus
 Proventriculus
 Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. Large intestine
- 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 7-3, E-F
Type of Bird: Senegal Parrot
Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 7 ml

administered via gavage tube Projection: Laterolateral (right lateral recumbency)

Weight of Animal: 146 g Gender: Unknown

Reproductive Status: Intact Age: Adult

Image	Time (hr)
Е	3.5
F	5.5

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus 5. Intestines
- 6. Duodenum
- 7. Large intestine
- 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 7-3, GType of Bird: Senegal Parrot
Type of Study: Gastrointestinal positive contrast study Contrast Study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 7 ml administered via gavage tube Projection: Laterolateral (right lateral recumbency) Weight of Animal: 146 g Gender: Unknown Reproductive Status: Intact Age: Adult

Image	Time (hr)
G	7.5

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. Large intestine
- 8. Cloaca
- 9. Proventricular-ventricular isthmus




Figure 7-4, A-BType of Bird: Senegal Parrot
Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)
7 ml administered via gavage tube

Projection: Ventrodorsal Weight of Animal: 146 g Gender: Unknown Reproductive Status: Intact Age: Adult

lmage	Time (hr)
A	Scout
R	0.5

- Crop
 (Esophagus)
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
- 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 7-4, C-DType of Bird: Senegal Parrot
Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)
7 ml administered via gavage tube

Projection: Ventrodorsal Weight of Animal: 146 g Gender: Unknown Reproductive Status: Intact Age: Adult

lmage	Time (hr)
С	1.5
D	2.5

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus5. Intestines
- 6. Duodenum
- 7. (Large intestine)
- 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 7-4, E-F

Type of Bird: Senegal Parrot
Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)
7 ml administered via gavage tube

Projection: Ventrodorsal Weight of Animal: 146 g Gender: Unknown Reproductive Status: Intact

Age: Adult

lmage	Time (hr)
Е	3.5
F	5.5

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)
- 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 7-4, GType of Bird: Senegal Parrot
Type of Study: Gastrointestinal positive

contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)
7 ml administered via gavage tube

Projection: Ventrodorsal Weight of Animal: 146 g Gender: Unknown Reproductive Status: Intact

Age: Adult

lmage	Time (hr)
G	7.5

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 7-5, A-B

Type of Bird: Senegal Parrot

Type of Study: Gastrointestinal double

contrast study. Gasdrollitestinal double contrast study. Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 7 ml and room air 7 ml administered

via gavage tube Projection: Laterolateral (right lateral recumbency)

Weight of Animal: 146 g Gender: Unknown Reproductive Status: Intact Age: Adult

Image	Time (hr)
A	Scout
В	0.25

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 7-5, CType of Bird: Senegal Parrot

Type of Study: Gastrointestinal double contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 7 ml and room air 7 ml administered via gavage tube

Projection: Laterolateral (right lateral

recumbency)
Weight of Animal: 146 g Gender: Unknown Reproductive Status: Intact

Age: Adult

lmage	Time (hr)
С	2.0

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus
- 5. Intestines6. (Duodenum)
- 7. Large intestine
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 7-6, A-B

Type of Bird: Senegal Parrot
Type of Study: Gastrointestinal double

contrast study
Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)
7 ml and room air 7 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 146 g Gender: Unknown Reproductive Status: Intact Age: Adult

Image	Time (hr)
A	Scout
В	0.25

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus
 5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
 8. Cloaca
- 9. Proventricular-ventricular isthmus




Figure 7-6, C
Type of Bird: Senegal Parrot
Type of Study: Gastrointestinal double contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 7 ml and room air 7 ml administered via gavage tube


Projection: Ventrodorsal Weight of Bird: 146 g Gender: Unknown Reproductive Status: Intact

Age: Adult

lmage	Time (hr)
С	2.0

- Crop
 (Esophagus)
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. Large intestine 8. Cloaca
- 9. (Proventricular-ventricular isthmus)

African Grey Parrot (Psittacus erithacus)






Figure 8-1, A-B Type of Bird: African Grey Parrot

Type of Study: Viscera of the coelom
Projection: Laterolateral (right lateral recumbency)

Weight of Animal: 546 g Gender: Unknown

Reproductive Status: Intact

Age: Adult

- 1. Trachea
- 2. Crop
- 3. Brachiocephalic artery and aorta
- 4. (Brachiocephalic artery)
- 5. (Aorta)6. Pulmonary artery
- 7. Pulmonary vein
- 8. Heart
- 9. Left atrium
- 10. Esophagus
- 11. Liver
- 12. Lung
- 13. Syrinx
- 14. Gonad 15. (Ovary) 16. (Testes)

- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. Cloaca
- 21. Cervical air sac
- 22. (Clavicular air sac)
- 23. Thoracic air sac
- 24. Abdominal air sac 25. Apex of heart
- 26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys
- 28. Spleen




Figure 8-2, A-B Type of Bird: African Grey Parrot
Type of Study: Viscera of the coelom
Projection: Ventrodorsal
Weight of Animal: 546 g Gender: Unknown Reproductive Status: Intact Age: Adult

- 1. Trachea

- 2. Cervical air sac
 3. Crop
 4. Clavicular air sac
 5. Brachiocephalic artery and aorta

 6. Grand 1. December 2. De
- 6. (Heart base vessel)
- 7. Heart
- 8. Lung
 9. Thoracic air sac

- 10. Liver
- 11. Kidneys12. Proventriculus
- 13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. (Cloaca)




Figure 8-3, A-B

Type of Bird: African Grey Parrot Type of Study: Whole body skeleton Projection: Laterolateral (right lateral recumbency) Weight of Animal: 546 g

Gender: Unknown Reproductive Status: Intact Age: Adult

- 3. Humerus 4. Coracoid
- 5. Scapula
- 6. Rib
 7. Uncinate process of rib
 8. Thoracic vertebra
- 9. Synsacrum 10. Ilium
- 11. Head of femur

- 13. Ischium
- 14. Pubis
- 15. Femur
- 16. Obturator foramen
- 17. Patella
- 18. Tibiotarsal bone
- 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae
- 22. Pygostyle




Figure 8-4, A-B Type of Bird: African Grey Parrot Type of Study: Whole body skeleton Projection: Ventrodorsal Weight of Animal: 546 g Gender: Unknown Reproductive Status: Intact Age: Adult

- 1. Cervical vertebra
- 2. Clavicle
- 3. Humerus
- 4. Coracoid 5. Scapula 6. Rib

- 7. Uncinate process of rib
- 8. Thoracic vertebra
- 9. Synsacrum
- 10. Ilium
- 11. Head of femur
- 12. (Ilioischiadiac foramen)
- 13. İschium
- 14. Pubis
- 15. Femur

- 16. (Obturator foramen)
- 17. (Patella)
- 18. Tibiotarsal bone
- 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae
- 22. Pygostyle
- 23. (Apex carinae)24. Dorsal tubercle of humerus25. Head of humerus
- 26. Ventral tubercle of humerus
- 27. Trochanter of femur




Figure 8-5
Type of Bird: African Grey Parrot
Type of Study: Infraorbital sinus contrast study

Contrast Medium: Iopamidol injection 41% (Isovue®-200, Bracco Diagnostics Inc., Princeton, NJ 08543) 6 ml instilled into the right nare

Projection: Laterolateral (right lateral

recumbency)
Weight of Animal: 422 g

Gender: Male Reproductive Status: Intact

Age: 2 years

- 1. Choana
- 2. External nares
- 3. Infraorbital sinus

- 4. Contrast medium in pharynx5. Endotracheal tube




Figure 8-6

Type of Bird: African Grey Parrot
Type of Study: Infraorbital sinus contrast
study

Contrast Medium: Iopamidol injection 41% (Isovue®-200, Bracco Diagnostics Inc., Princeton, NJ 08543) 6 ml instilled into the right nare Projection: Ventrodorsal Weight of Animal: 422 g


Gender: Male

Reproductive Status: Intact Age: 2 years

- 1. Choana
- 2. External nare3. Infraorbital sinus

- 4. Contrast medium in pharynx
- 5. Endotracheal tube

Orange-Winged Amazon Parrot (Amazona amazonica)






Figure 9-1, A-B

Type of Bird: Orange-Winged Amazon

Parrot

Body Condition: Normal

Type of Study: Viscera of the coelom

Projection: Laterolateral (right lateral

recumbency)
Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact

Age: 13 years

- 1. Trachea
- 2. Crop
- 3. Brachiocephalic artery and aorta
- 4. (Brachiocephalic artery)
- 5. Aorta
- 6. Pulmonary artery
- 7. Pulmonary vein
- 8. Heart
- 9. Left atrium
- 10. Esophagus
- 11. Liver
- 12. Lung
- 13. Syrinx 14. Gonad
- 15. (Ovary) 16. (Testes)

- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. Cloaca
- 21. (Cervical air sac)
- 22. (Clavicular air sac)
- 23. (Thoracic air sac)
- 24. Abdominal air sac 25. (Apex of heart)
- 26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys
- 28. Spleen




Figure 9-2, A-B
Type of Bird: Orange-Winged Amazon
Parrot Body Condition: Normal Type of Study: Viscera of the coelom Projection: Ventrodorsal Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact Age: 13 years

- Trachea
 Cervical air sac
 Crop
- 4. Clavicular air sac
- 5. Brachiocephalic artery and aorta
- 6. Heart base vessel 7. Heart
- 8. Lung
- 9. Thoracic air sac
- 10. Liver

- 11. Kidneys12. Proventriculus
- 13. (Ventriculus)
- 14. Intestines
- 15. Abdominal air sac
- 16. Cloaca




Figure 9-3, A-BType of Bird: Orange-Winged Amazon Parrot

Body Condition: Obese

Type of Study: Viscera of the coelom Projection: Laterolateral (right lateral

recumbency) Weight of Bird: 580 g Gender: Unknown

Reproductive Status: Intact Age: Adult

- 1. Trachea
- 2. Crop3. Brachiocephalic artery and aorta
- 4. (Brachiocephalic artery)
- 5. (Aorta)
- 6. Pulmonary artery
- 7. Pulmonary vein8. Heart
- 9. Left atrium
- 10. Esophagus
- 11. Liver
- 12. Lung
- 13. Syrinx 14. Gonad
- 15. (Ovary) 16. (Testes)

- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. Cloaca
- 21. Cervical air sac
- 22. Clavicular air sac
- 23. Thoracic air sac
- 24. Abdominal air sac
- 25. Apex of heart
- 26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys
- 28. Spleen




Figure 9-4, A-B
Type of Bird: Orange-Winged Amazon
Parrot
Parrot

Body Condition: Obese Type of Study: Viscera of the coelom Projection: Ventrodorsal Weight of Bird: 580 g Gender: Unknown

Reproductive Status: Intact

Age: Adult

- 1. Trachea
- 2. (Cervical air sac)3. Crop
- 4. Clavicular air sac
- 5. Brachiocephalic artery and aorta
- 6. Heart base vessel
- 7. Heart
- 8. Lung
- 9. Thoracic air sac
- 10. Liver

- 11. Kidneys12. Proventriculus
- 13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. Cloaca




Figure 9-5, A-B

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: Head Projection: Laterolateral (right lateral

recumbency) Weight of Bird: 412 g

Gender: Unknown Reproductive Status: Intact

Age: 13 years

- 1. Keratinized maxillary beak
- 2. Keratinized mandibular beak
- 3. Premaxillary bone
- 4. Mandible
- 5. Hyoid bones
- 6. Palatine bone
- 7. Pterygoid bone

- 8. External nares
 9. Jugal [zygomatic] bone
 10. Craniofacial flexion zone
- 11. Orbit
- 12. Frontal bone
- 13. Cranium
- 14. Temporal bone 15. (Quadrate bone)
- 16. Parietal bone




Figure 9-6, A-B

Type of Bird: Orange-Winged Amazon
Parrot

Type of Study: Head
Projection: Oblique 30% Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact

Age: 13 years

- 1. Keratinized maxillary beak
- 2. Keratinized mandibular beak
- 3. Premaxillary bone4. Mandible
- 5. Hyoid bones
- 6. Palatine bone
 7. (Pterygoid bone)
- 8. External nares
- 9. Jugal [zygomatic] bone
- 10. Craniofacial flexion zone

- 11. Orbit
- 12. Frontal bone
- 13. Cranium
- 14. Temporal bone
- 15. (Quadrate bone)
- 16. Parietal bone



Figure 9-7, A

Type of Bird: Orange-Winged Amazon
Parrot
Type of Study: Head
Projection: Ventrodorsal
Weight of Bird: 412 g
Gender: Unknown
Reproductive Status: Intact
Age: 13 years




Figure 9-7, B

Type of Bird: Orange-Winged Amazon
Parrot
Type of Study: Head
Projection: Ventrodorsal
Weight of Bird: 412 g
Gender: Unknown
Reproductive Status: Intact
Age: 13 years

Age: 13 years

- 1. Keratinized maxillary beak
- Relatilized maxilia
 Premaxillary bone
 Mandible
 Orbit

- 5. Jugal [zygomatic] bone6. Scleral ossicles7. Quadrate bone8. Cranium




Figure 9-8, A-BType of Bird: Orange-Winged Amazon Parrot

Type of Study: Whole body skeleton Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 412 g

Gender: Unknown Reproductive Status: Intact

Age: 13 years

- Cervical vertebra
 Clavicle
 Humerus
- 4. Coracoid
- 5. Scapula
- 6. Rib
 7. (Uncinate process of rib)
- 8. Thoracic vertebra
- 9. Synsacrum
- 10. Ilium
- 11. (Head of femur)
- 12. Ilioischiadiac foramen
- 13. Ischium

- 14. Pubis
- 15. Femur
- 16. Obturator foramen
- 17. Patella
- 18. Tibiotarsal bone
- 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae22. Pygostyle




Figure 9-9, A-B
Type of Bird: Orange-Winged Amazon
Parrot Type of Study: Whole body skeleton Projection: Ventrodorsal Weight of Bird: 412 g Gender: Unknown

Reproductive Status: Intact

Age: 13 years

- 1. Cervical vertebra
- Clavicle
 Humerus
- Coracoid
- 5. Scapula6. Rib7. (Uncinate process of rib)
- 8. Thoracic vertebra
- 9. Synsacrum
- 10. Ilium
- 11. Head of femur
- 12. (Ilioischiadiac foramen)
- 13. Ischium
- 14. Pubis
- 15. Femur

- 16. (Obturator foramen)
 17. Patella
- 18. Tibiotarsal bone
- 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae
- 22. Pygostyle
- 23. Apex carinae
- 24. Dorsal tubercle of humerus
- 25. Head of humerus
- 26. Ventral tubercle of humerus
- 27. Trochanter of femur




Figure 9-10, A-B

Type of Bird: Orange-Winged Amazon
Parrot
Parrot

Type of Study: Wing Projection: Mediolateral Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact

Age: 13 years

- 1. Head of humerus
- Dorsal tubercle of humerus
 Dorsal condyle of humerus
- 4. Radius
- 5. Radial carpal bone
- 6. Alula
- 7. Major metacarpal bone
- 8. Phalanges of major digit
- 9. Ventral tubercle of humerus
- 10. Humerus
 11. Ventral condyle of humerus
- 12. Ulna
- 13. Ulnar carpal bone
- 14. Minor metacarpal bone 15. Phalanges of minor digit




Figure 9-11, A-B
Type of Bird: Orange-Winged Amazon
Parrot

Type of Study: Wing Projection: Caudocranial Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact

Age: 13 years

- Head of humerus
 Ventral tubercle of humerus
 Ventral condyle of humerus
- 4. Radius

- 5. Radial carpal bone6. Alula7. (Minor metacarpal bone)
- 8. Phalanges of major digit9. Dorsal tubercle of humerus
- 10. Humerus
- 11. Dorsal condyle of humerus
- 12. Ulna
- 13. (Ulnar carpal bone)
- 14. Major metacarpal bone 15. (Phalanges of minor digit)




Figure 9-12, A-B
Type of Bird: Orange-Winged Amazon Parrot Type of Study: Pelvic limb Projection: Mediolateral Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact Age: 13 years

- 1. Femur
- 2. Patella
- 3. Condyles of femur
 4. Fibula
- 5. Tibiotarsal bone
- 6. Condyles of tibiotarsal bone
- 7. Hypotarsal crest of tarsometatarsal bone
- 8. Tarsometatarsal bone

- 9. Trochlea of tarsometatarsal bone
- 10. Digit I

- 11. Digit II 12. Digit III 13. Digit IV
- 14. Phalanges
- 15. Metatarsal I




Figure 9-13, A-B

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: Pelvic limb Projection: Craniocaudal Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact

Age: 13 years

- 1. Trochanter of femur
- 2. Femur3. Lateral condyle of femur
- 4. Fibula
- 5. Lateral condyle of tibiotarsal bone6. Tarsometatarsal bone
- 7. Trochlea of tarsometatarsal bone 8. Digit IV
- 9. Digit III
- 10. Head of femur
- 11. Medial condyle of femur12. Tibiotarsal bone

- 13. Medial condyle of tibiotarsal bone14. Digit I15. Digit II

- 16. Phalanges
- 17. (Patella)
- 18. Hypotarsal crest of tarsometatarsal bone
- 19. Metatarsal I




Figure 9-14, A-BType of Bird: Orange-Winged Amazon Parrot Type of Study: Distal pelvic limb Projection: Mediolateral Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact

Age: 13 years

- Tarsometatarsal bone
 Trochlea of tarsometatarsal bone
 Metatarsal bone I

- 4. Digit I 5. Digit II 6. Digit III

- 7. Digit IV8. Phalanges
- 9. Hypotarsal crest of tarsometatarsal bone
- 10. Condyles of tibiotarsal bone




Figure 9-15, A-B

Type of Bird: Orange-Winged Amazon
Parrot

P Type of Study: Distal pelvic limb Projection: Dorsoplantar Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact Age: 13 years

- Tarsometatarsal bone
 Trochlea of tarsometatarsal bone
 Metatarsal bone I

- 4. Digit I 5. Digit II 6. Digit III

- 7. Digit IV 8. Phalanges 9. Hypotarsal crest of tarsometatarsal
- 10. Condyles of tibiotarsal bone




Figure 9-16, A-B

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)

15 ml administered via gavage tube Projection: Laterolateral (right lateral recumbency) Weight of Bird: 412 g

Gender: Unknown Reproductive Status: Intact

Age: 13 years

lmage	Time (hr)
A	Scout
В	0.25

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
 7. (Large intestine)
 8. Cloaca
- 9. Proventricular-ventricular isthmus




Figure 9-16, C-DType of Bird: Orange-Winged Amazon Parrot

Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 15 ml administered via gavage tube

Projection: Laterolateral (right lateral

recumbency)
Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact

Age: 13 years

lmage	Time (hr)
С	0.5
D	1.0

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)
- 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 9-16, E-F

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 15 ml administered via gavage tube Projection: Laterolateral (right lateral

recumbency)

Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact

Age: 13 years

Image	Time (hr)
E	2.0
F	4.0

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)
- 8. Cloaca
- 9. Proventricular-ventricular isthmus




Figure 9-17, A-BType of Bird: Orange-Winged Amazon Parrot

Type of Study: Gastrointestinal positive

contrast study
Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 15 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact Age: 13 years

lmage	Time (hr)
A	Scout
В	0.25

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
 6. (Duodenum)
- 7. (Large intestine)
- 8. Cloaca
- 9. Proventricular-ventricular isthmus




Figure 9-17, C-D

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 15 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact Age: 13 years

Time (hr) Image C 0.5 D 1.0

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)
 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 9-17, E-F
Type of Bird: Orange-Winged Amazon
Parrot

Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 15 ml administered via gavage tube Projection: Ventrodorsal Weight of Bird: 412 g

Gender: Unknown Reproductive Status: Intact

Age: 13 years

lmage	Time (hr)
Е	2.0
F	4.0

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
- 8. Cloaca
- 9. Proventricular-ventricular isthmus




Figure 9-18, A-B

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: Gastrointestinal double contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 10 ml and room air 60 ml administered via gavage tube

Projection: Laterolateral (right lateral

recumbency)
Weight of Bird: 360 g Gender: Male

Reproductive Status: Intact

Age: 15 years

lmage	Time (hr)
A	Scout
В	0.083

- 1. Crop
- Esophagus
 Proventriculus
 Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
- 8. Cloaca
- $9. \ (Proven tricular-ventricular\ is thmus)$




Figure 9-18, C-D

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: Gastrointestinal double contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 10 ml and room air 60 ml administered

via gavage tube Projection: Laterolateral (right lateral

recumbency) Weight of Bird: 360 g

Gender: Male Reproductive Status: Intact

Age: 15 years

lmage	Time (hr)
С	0.25
D	1.0

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines6. Duodenum
- 7. Large intestine
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 9-19, A-B

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: Gastrointestinal double contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 10 ml and room air 60 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: $360\ g$ Gender: Male

Reproductive Status: Intact

Age: 15 years

Image	Time (hr)
A	Scout
В	0.083

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
 8. (Cloaca)
- $9. \ Proven tricular-ventricular \ is thmus$




Figure 9-19, C-D
Type of Bird: Orange-Winged Amazon
Parrot

Type of Study: Gastrointestinal double contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 10 ml and room air 60 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 360 g Gender: Male

Reproductive Status: Intact

Age: 15 years

lmage	Time (hr)
С	0.25
D	1.0

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus
- $5. \ Intestines$
- 6. (Duodenum)
- 7. (Large intestine)
- 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 9-20, A-BType of Bird: Orange-Winged Amazon Parrot

Type of Study: Cloacagram (retrograde, positive and double contrast)

Contrast Medium: Iopamidol injection 41% (Isovue®-200, Bracco Diagnostics Inc.,

Princeton, NJ 08543) 1.8 ml (positive contrast study); followed by room air 3 ml via vent (double contrast study) after partial removal of contrast medium

Projection: Laterolateral (right lateral recumbency)

Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact

Age: 13 years

Image	Study Type
A	Scout
В	Retrograde Positive Contrast Study

- 1. Colon
- Coprodeum
 Urodeum
- 4. Cloaca
- 5. Vent6. Intestines




Figure 9-20, C

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: Cloacagram (retrograde, positive and double contrast)

Contrast Medium: Iopamidol injection 41% (Isovue®-200, Bracco Diagnostics Inc., Princeton, NJ 08543) 1.8 ml (positive contrast study); followed by room air 3 ml via vent (double contrast study) after partial removal of contrast medium

Projection: Laterolateral (right lateral

recumbency) Weight of Bird: 412 g Gender: Unknown

Reproductive Status: Intact

Age: 13 years

lmage	Study Type
С	Double Contrast Study

- 1. Colon
- 2. (Coprodeum)
 3. (Urodeum)
 4. Cloaca
 5. (Vent)

- 6. Intestines




Figure 9-21, A-B

Type of Bird: Orange-Winged Amazon
Parrot

Type of Study: Cloacagram (retrograde, positive and double contrast)
Contrast Medium: Iopamidol injection 41%

(Isovue®-200, Bracco Diagnostics Inc., Princeton, NJ 08543) 1.8 ml (positive contrast study) followed by room air 3 ml via vent (double contrast study) after partial removal of contrast medium

Projection: VentrodorsalWeight of Bird: 412 g Gender: Unknown Reproductive Status: Intact Age: 13 years

Image Study Type A Scout Retrograde Positive В Contrast Study

- 1. (Colon)
 2. (Coprodeum)
 3. (Urodeum)

- 4. Cloaca 5. (Vent)
- 6. Intestines



Figure 9-21, C

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: Cloacagram (retrograde, positive and double contrast)

Contrast Medium: Iopamidol injection 41% (Isovue®-200, Bracco Diagnostics Inc., Princeton, NJ 08543) 1.8 ml (positive contrast study) followed by room air 3 ml via vent (double contrast study) after partial removal of contrast medium

Projection: Ventrodorsal Weight of Bird: 412 g Gender: Unknown Reproductive Status: Intact Age: 13 years ImageStudy TypeCDouble Contrast
Study

- 1. (Colon)
- 2. (Coprodeum)
- 3. (Urodeum)
- 4. Cloaca
- 5. Vent
- 6. Intestines






Figure 9-22, A-B
Type of Bird: Orange-Winged Amazon
Parrot
Parrot

Type of Study: Excretory urogram Contrast Medium: Diatrizoate meglumine, diatrizoate sodium (RenoCal – 76® 37% organically bound iodine; Bracco Diagnostics, Inc., Princeton, NJ) 0.72 ml

IV (2 ml/kg) via cutaneous ulnar vein Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 360 g

Gender: Male

Reproductive Status: Intact

Age: 15 years

lmage	Time (min)
A	Scout
В	10

- Kidneys
 Ureter
 Cloaca



Figure 9-22, C

Type of Bird: Orange-Winged Amazon
Parrot

Type of Study: Excretory urogram

Contrast Medium: Diatrizoate meglumine,
diatrizoate sodium (RenoCal-76® 37%
organically bound inding: Bracco Diag organically bound iodine; Bracco Diagnostics, Inc., Princeton, NJ) 0.72 ml IV (2 ml/kg) via cutaneous ulnar vein Projection: Laterolateral (right lateral

recumbency)

Weight of Animal: 360 g

Gender: Male

Reproductive Status: Intact Age: 15 years

Image	Time (min)
С	15

- Kidneys
 Ureter
- 3. Cloaca




Figure 9-23, A-B
Type of Bird: Orange-Winged Amazon
Parrot
Parrot

Type of Study: Excretory urogram

Contrast Medium: Diatrizoate meglumine, diatrizoate sodium (RenoCal–76® 37% organically bound iodine; Bracco Diagnostics, Inc., Princeton, NJ) 0.72 ml IV (2 ml/kg) via cutaneous ulnar vein

Projection: Ventrodorsal Weight of Bird: 360 g

Gender: Male Reproductive Status: Intact

Age: 15 years

Time (min) Image A Scout В 10

- Kidneys
 Ureter
 Cloaca



Figure 9-23, C

Type of Bird: Orange-Winged Amazon
Parrot
Type of Study: Excretory urogram Contrast Medium: Diatrizoate meglumine, diatrizoate sodium (RenoCal–76® 37% organically bound iodine; Bracco Diagnostics, Inc., Princeton, NJ) 0.72 ml IV (2 ml/kg) via cutaneous ulnar vein

Projection: Ventrodorsal Weight of Bird: 360 g

Gender: Male Reproductive Status: Intact

Age: 15 years

lmage	Time (min)
С	15

- Kidneys
 Ureter
 Cloaca




Figure 9-24, A-D

Type of Bird: Orange-Winged Parrot Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 412 kg Age: 13 years

- Tongue
 Mandible
- 3. (Hyoid bone)
- 4. Endotracheal tube
- 5. Keratinized maxillary beak6. (Nasal cavity)7. Nasal concha

- 8. Pharynx
- 9. Premaxillary bone
- 10. (Infraorbital sinus)
- 11. Nasal septum
- 12. (Choana)
- 13. Palatine bone
- 14. (Sphenoid bone) 15. (Jugal [zygomatic] bone)
- 16. (Frontal bone)
- 17. (Pterygoid bone) 18. (Eyeball)

- 19. (Scleral ossicle)
- 20. (Interorbital septum)21. (Lens of eyeball)

- 22. (Trachea)
 23. (Cerebrum)
- 24. (External ear canal)
- 25. (Cerebellum)
- 26. (Spinal cord)
- 27. (Dens) 28. (Cere)
- 29. (Nare[s])
- 30. (Feather)
- 31. (Pons)
- 32. (Occipital bone)
 33. (Cervical vertebra)




Figure 9-24, E-H

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 412 kg Age: 13 years

- 1. Tongue
- 2. Mandible
- 3. Hyoid bone
- 4. (Endotracheal tube)
- 5. (Keratinized maxillary beak)
- 6. Nasal cavity
- 7. Nasal concha
- 8. Pharynx
- 9. (Premaxillary bone)
 10. Infraorbital sinus
- 11. (Nasal septum)
- 12. Choana
- 13. Palatine bone 14. (Sphenoid bone)
- 15. Jugal [zygomatic] bone
- 16. (Frontal bone)
- 17. (Pterygoid bone)
- 18. Eyeball

- 19. Scleral ossicle
- 20. Interorbital septum 21. (Lens of eyeball)

- 22. (Trachea)
 23. (Cerebrum)
- 24. (External ear canal)
- 25. (Cerebellum)
- 26. (Spinal cord) 27. (Dens)
- 28. (Cere)
- 29. (Nare[s])
- 30. (Feather)
- 31. (Pons)
- 32. (Occipital bone) 33. (Cervical vertebra)




Figure 9-24, I-KType of Bird: Orange-Winged Amazon Parrot Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 412 kg Age: 13 years

5. (Keratinized maxillary beak) 6. (Nasal cavity)
7. (Nasal concha) 8. (Pharynx) 9. (Premaxillary bone) 10. (Infraorbital sinus) 11. (Nasal septum)
12. (Choana)
13. Palatine bone 14. (Sphenoid bone)

3. Hyoid bone

18. (Eyeball)

4. (Endotracheal tube)

- 15. Jugal [zygomatic] bone 16. Frontal bone 17. Pterygoid bone
- 24. (External ear canal)
 25. (Cerebellum) 26. (Spinal cord) 27. (Dens) 28. (Cere) 29. (Nare[s]) 30. (Feather) 31. (Pons) 32. (Occipital bone) 33. (Cervical vertebra)

21. Lens of eyeball

22. (Trachea) 23. Cerebrum




Figure 9-25, A-D
Type of Bird: Orange-Winged Amazon Parrot

Type of Study: CT coelom Contrast Medium: None Imaging Plane: Transverse Weight of bird: 412 kg Age: 13 years

- 1. Sternum
- 2. Thoracic air sac
- 3. (Lung)
- 4. (Esophagus)
 5. Thoracic vertebra
 6. (Heart)
 7. (Blood vessels)

- 7. (Blood vessels)8. Thoracic extremity [wing]
- 9. (Proventriculus)
 10. (Ventriculus)
- 11. (Intestines)
- 12. (Pelvic extremity [leg])
- 13. (Abdominal air sac) 14. (Ilium) 15. (Liver)

- 16. (Kidney) 17. (Caudal vertebra)
- 18. (Rib)
- 19. (Aorta)
 20. (Pubic bone)
 21. (Cloaca)
- 22. (Clavicular air sac)23. Scapula
- 24. Syrinx
- 25. (Ischium) 26. (Colon)
- 27. (Spinal cord)




Figure 9-25, E-HType of Bird: Orange-Winged Amazon Parrot Type of Study: CT coelom

Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 412 kg

Age: 13 years

- (Sternum)
 Thoracic air sac
- 3. (Lung) 4. Esophagus
- 5. Thoracic vertebra
- 6. Heart7. Blood vessels
- 8. (Thoracic extremity [wing])
- 9. Proventriculus
- 10. Ventriculus
- 11. Intestines
- 12. Pelvic extremity [leg]
- 13. Abdominal air sac
- 14. Ilium
- 15. Liver

- 16. Kidney
- 17. (Caudal vertebra)
- 18. Rib
- 19. (Aorta)
 20. Pubic bone
 21. (Cloaca)
- 22. (Clavicular air sac)
- 23. (Scapula)
- 24. (Syrinx) 25. Ischium 26. (Colon)
- 27. (Spinal cord)




Figure 9-25, I-J
Type of Bird: Orange-Winged Amazon
Parrot

Type of Study: CT coelom Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 412 kg

Age: 13 years

- 1. (Sternum)
 2. (Thoracic air sac)
 3. (Lung)

- 4. (Esophagus)5. (Thoracic vertebra)
- 6. (Heart)
- 7. (Blood vessels)
 8. (Thoracic extremity [wing])
- 9. (Proventriculus)
- 10. (Ventriculus)
- 11. Intestines
 12. Pelvic extremity [leg]
 13. (Abdominal air sac)
- 14. (Ilium)
- 15. (Liver)

- 16. (Kidney) 17. Caudal vertebra 18. (Rib)
- 19. (Aorta)
- 20. Pubic bone
- 21. Cloaca
- 22. (Clavicular air sac)
- 23. (Scapula)
- 24. (Syrinx)
- 25. Ischium
- 26. Colon
- 27. (Spinal cord)




Figure 9-26, A-DType of Bird: Orange-Winged Amazon Parrot

Type of Study: CT coelom (post-contrast)
Contrast Medium: Iopamidol injection
41% (Isovue®-200, Bracco Diagnostics Inc., Princeton, NJ 08543) 0.4 ml (1 ml/kg) intravenously via cutaneous ulnar vein

Imaging Plane: Transverse Weight of Bird: 412 kg Age: 13 years

- 1. Sternum
- 2. Thoracic air sac
- 3. (Lung)
- 4. Esophagus
- 5. (Thoracic vertebra)
- 6. Heart
- (Blood vessels)
- 8. (Thoracic extremity [wing])
- 9. Proventriculus
- 10. (Ventriculus)
- 11. (Intestines)
- 12. (Pelvic extremity [leg])
- 13. (Abdominal air sac)
- 14. İlium
- 15. Liver

- 16. (Kidney) 17. (Caudal vertebra)
- 18. Rib

- 19. (Aorta)
 20. (Pubic bone)
 21. (Cloaca)
 22. (Clavicular air sac)
- 23. (Scapula)
- 24. (Syrinx)
- 25. (Ischium)
- 26. (Colon) 27. (Spinal cord)




Figure 9-26, E-GType of Bird: Orange-Winged Amazon Parrot

Type of Study: CT coelom (post-contrast) Contrast Medium: Iopamidol injection 41% (Isovue®-200, Bracco Diagnostics Inc., Princeton, NJ 08543) 0.4 ml (1 ml/kg) intravenously via cutaneous ulnar vein

Imaging Plane: Transverse Weight of Bird: 412 kg Age: 13 years

- 1. (Sternum)
- 2. (Thoracic air sac)
- 3. (Lung)
- 4. (Esophagus)
- 5. (Thoracic vertebra)
- 6. (Heart)
 7. Blood vessels
- 8. (Thoracic extremity [wing])
- 9. (Proventriculus)
- 10. Ventriculus
- 11. Intestines
- 12. Pelvic extremity [leg]
- 13. (Abdominal air sac)
- 14. İlium
- 15. (Liver)

- 16. Kidney
- 17. Caudal vertebra
- 18. (Rib)
- 19. (Aorta)
- 20. Pubic bone
- 21. Cloaca
- 22. (Clavicular air sac)
- 23. (Scapula)
- 24. (Syrinx) 25. (Ischium) 26. (Colon)
- 27. (Spinal cord)




Figure 9-27, A-B

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: MRI head (T1) Contrast Medium: None Imaging Plane: Sagittal Weight of Bird: 360 g Age: 15 years

1. (Tongue)

2. (Mandible)

3. Hyoid bone
4. (Endotracheal tube)

5. (Keratinized maxillary beak)

6. (Nasal cavity) (Nasal concha)

8. (Pharynx)

9. (Premaxillary bone)

10. (Infraorbital sinus)

11. (Nasal septum)

12. (Choana)
13. (Palatine bone)

14. (Sphenoid bone)

15. (Jugal [zygomatic] bone)

16. (Frontal bone)

17. (Pterygoid bone)

18. (Eyeball)

19. (Scleral ossicle)

20. (Interorbital septum)

21. (Lens of eyeball)

22. Trachea

23. Cerebrum

24. (External ear canal)

25. Čerebellum 26. Spinal cord

27. (Dens)

28. (Cere)

29. (Nare[s])

30. (Feather)

31. (Pons)

32. (Occipital bone)

33. (Cervical vertebra)




Figure 9-27, C-D

Type of Bird: Orange-Winged Amazon Parrot Type of Study: MRI head (T1)

Contrast Medium: None Imaging Plane: Sagittal Weight of Bird: 360 g Age: 15 years

2. (Mandible) 3. (Hyoid bone)
4. (Endotracheal tube)

1. (Tongue)

5. (Keratinized maxillary beak)

6. (Nasal cavity) (Nasal concha) 8. (Pharynx)

9. (Premaxillary bone) 10. (Infraorbital sinus)

11. (Nasal septum) 12. (Choana)

13. (Palatine bone) 14. (Sphenoid bone)

15. (Jugal [zygomatic] bone)

16. (Frontal bone) 17. (Pterygoid bone)

18. Eyeball

19. (Scleral ossicle)

20. (Interorbital septum)

21. Lens of eyeball

22. (Trachea)

23. Cerebrum

24. (External ear canal)

25. (Cerebellum)

26. (Spinal cord) 27. (Dens)

28. (Cere)

29. (Nare[s]) 30. (Feather)

31. (Pons)

32. (Occipital bone)

33. (Cervical vertebra)




Figure 9-28, A-C

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: MRI head (T1 postcontrast)

Contrast Medium: Gadopentetate dimeglumine (Magnevist® injection, Bayer Health Care Pharmaceuticals, Wayne NJ 07470) 0.07 ml (0.2 ml/kg) administered directly into a catheter in the cutaneous ulnar vein.

Imaging Plane: Sagittal Weight of Bird: 360 g Age: 15 years

- (Tongue)
 (Mandible)
- 3. (Hyoid bone)
- 4. (Endotracheal tube)
- 5. (Keratinized maxillary beak)
- 6. (Nasal cavity)
- (Nasal concha)
- 8. Pharynx
- 9. (Premaxillary bone)
- 10. (Infraorbital sinus)
- 11. (Nasal septum)
- 12. (Choana)
- 13. (Palatine bone) 14. (Sphenoid bone)
- 15. (Jugal [zygomatic] bone)
- 16. (Frontal bone)
- 17. (Pterygoid bone)
- 18. Eyeball

- 19. (Scleral ossicle)
- 20. (Interorbital septum)
- 21. Lens of eyeball
- 22. Trachea
- 23. Cerebrum
- 24. (External ear canal)
- 25. Cerebellum
- 26. Spinal cord
- 27. (Dens)
- 28. (Cere)
- 29. (Nare[s])
- 30. (Feather) 31. (Pons)
- 32. (Occipital bone)
- 33. (Cervical vertebra)




Figure 9-29, A-CType of Bird: Orange-Winged Amazon Parrot Type of Study: MRI head (T1) Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 360 g Age: 15 years

- (Tongue)
 (Mandible)
- 3. (Hyoid bone)
- 4. (Endotracheal tube)
- 5. (Keratinized maxillary beak)
- 6. (Nasal cavity)
- 7. (Nasal concha)
- 8. Pharynx
- 9. (Premaxillary bone)
- 10. Infraorbital sinus
- 11. (Nasal septum)
- 12. (Choana)
- 13. (Palatine bone)
- 14. (Sphenoid bone)
- 15. (Jugal [zygomatic] bone)
- 16. (Frontal bone)
- 17. (Pterygoid bone)
- 18. Eyeball

- 19. (Scleral ossicle)
 20. (Interorbital septum)
- 21. Lens of eyeball
- 22. (Trachea)
- 23. Cerebrum
- 24. (External ear canal)
- 25. Cerebellum
- 26. (Spinal cord)
- 27. (Dens)
- 28. (Cere)
- 29. (Nare[s])
- 30. (Feather)
- 31. (Pons)
- 32. (Occipital bone)
- 33. (Cervical vertebra)




Figure 9-29, D-E

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: MRI head (T1) Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 360 g Age: 15 years

- (Tongue)
 (Mandible)
 (Hyoid bone)
- 4. (Endotracheal tube)
- 5. (Keratinized maxillary beak)
- 6. (Nasal cavity)
- (Nasal concha)
- 8. (Pharynx)
- 9. (Premaxillary bone)
- 10. (Infraorbital sinus)
- 11. (Nasal septum)
 12. (Choana)
 13. (Palatine bone)
- 14. (Sphenoid bone)
- 15. (Jugal [zygomatic] bone)
- 16. (Frontal bone)
 17. (Pterygoid bone)
- 18. (Eyeball)

- 19. (Scleral ossicle)20. (Interorbital septum)21. (Lens of eyeball)
- 22. (Trachea)
- 23. (Cerebrum)
- 24. (External ear canal)
 25. Cerebellum
- 26. Spinal cord
- 27. (Dens)
- 28. (Cere)
- 29. (Nare[s])
- 30. (Feather)
- 31. (Pons)
- 32. (Occipital bone)
- 33. (Cervical vertebra)




Figure 9-30, A-D

Type of Bird: Orange-Winged Amazon Parrot

Type of Study: MRI head (T1 post-contrast) Contrast Medium: Gadopentetate dimeglumine (Magnevist® injection, Bayer Health Care Pharmaceuticals, Wayne NJ 07470) 0.07 ml (0.2 ml/kg) administered directly into a catheter in the cutaneous ulnar vein.

Imaging Plane: Transverse Weight of Bird: 360 g Age: 15 years

- 1. (Tongue)
- 2. Mandible
- 3. (Hyoid bone)
- (Endotracheal tube)
- 5. (Keratinized maxillary beak)
- 6. (Nasal cavity)
- (Nasal concha) 7.
- 8. Pharynx
- 9. (Premaxillary bone)
- 10. Infraorbital sinus
- 11. (Nasal septum)
- 12. (Choana)
- 13. (Palatine bone)
- 14. (Sphenoid bone)
- 15. (Jugal [zygomatic] bone)
- 16. (Frontal bone)
- 17. (Pterygoid bone)
 18. Eyeball

- 19. (Scleral ossicle)
- 20. (Interorbital septum)
- 21. Lens of eyeball
- 22. (Trachea)
- 23. Cerebrum
- 24. (External ear canal)
- 25. Cerebellum
- 26. Spinal cord
- 27. (Dens)
- 28. (Cere)
- 29. (Nare[s])
- 30. (Feather)
- 31. (Pons)
- 32. (Occipital bone)
- 33. (Cervical vertebra)




Figure 9-31
Type of Bird: Orange-Winged Amazon
Parrot


Body Condition: Normal
Type of Study: Ultrasound study of the
liver

Projection: Sagittal Weight of Animal: 412 g Gender: Unknown Reproductive Status: Intact

Age: 13 years

- 1. Heart
- 2. Liver3. Intestines

Blue and Gold Macaw (Ara ararauna)






Figure 10-1, A-B

Type of Bird: Blue and Gold Macaw Type of Study: Viscera of the coelom Projection: Laterolateral (right lateral recumbency)

Weight of Bird: 1.1 kg Gender: Female

Reproductive Status: Intact Age: Adult

- 1. Trachea
- 2. Crop
- 3. Brachiocephalic artery and aorta
- 4. (Brachiocephalic artery)
- 5. (Aorta)
- 6. Pulmonary artery
- 7. Pulmonary vein
- 8. Heart
- 9. Left atrium
- 10. (Esophagus)
- 11. Liver
- 12. Lung
- 13. Syrinx 14. (Gonad) 15. (Ovary) 16. (Testes)

- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. Cloaca
- 21. Cervical air sac
- 22. (Clavicular air sac)
- 23. Thoracic air sac
- 24. Abdominal air sac 25. (Apex of heart)
- 26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys
- 28. Spleen




Figure 10-2, A-BType of Bird: Blue and Gold Macaw
Type of Study: Viscera of the coelom Projection: Ventrodorsal Weight of Bird: 1.1 kg Gender: Female Reproductive Status: Intact Age: Adult

- 1. Trachea
- 2. (Cervical air sac)3. Crop
- 4. Clavicular air sac
- 5. Brachiocephalic artery and aorta6. Heart base vessel7. Heart

- 8. Lung9. Thoracic air sac
- 10. Liver

- 11. Kidneys12. Proventriculus
- 13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. Cloaca




Figure 10-3, A-B
Type of Bird: Blue and Gold Macaw
Type of Study: Head

Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 1.1 kg

Gender: Female

Reproductive Status: Intact

Age: Adult

- 1. Keratinized maxillary beak
- 2. Keratinized mandibular beak
- 3. Premaxillary bone
- 4. Mandible
- 5. Hyoid bones 6. Palatine bone 7. Pterygoid bone
- 8. External nares

- 9. Jugal (zygomatic) bone
- 10. Craniofacial flexion zone
- 11. Orbit
- 12. Frontal bone

- 13. Cranium
 14. Temporal bone
 15. Quadrate bone
- 16. Parietal bone





Figure 10-4, A-B
Type of Bird: Blue and Gold Macaw
Type of Study: Head
Projection: Ventrodorsal
Weight of Bird: 1.1 kg
Gender: Female
Reproductive Status: Intact
Age: Adult

- Keratinized maxillary beak
 Premaxillary bone
 Mandible
 (Orbit)
 Jugal (zygomatic) bone
 Scleral ossicles

- 7. (Quadrate bone)8. Cranium



Type of Bird: Blue and Gold Macaw Type of Study: Whole body skeleton
Projection: Laterolateral (right lateral recumbency) Weight of Bird: 1.1 kg Gender: Female

Reproductive Status: Intact Age: Adult

- 3. Humerus
- 4. Coracoid
- 5. Scapula 6. Rib
- 7. (Uncinate process of rib)
- 8. Thoracic vertebra
- 9. Synsacrum
- 10. Ilium
- 11. Head of femur
- 12. Ilioischiadiac foramen
- 13. Ischium

- 16. Obturator foramen
- 17. Patella
- 18. Tibiotarsal bone
- 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae22. Pygostyle




Figure 10-6, A-B Type of Bird: Blue and Gold Macaw Type of Study: Whole body skeleton Projection: Ventrodorsal Weight of Bird: 1.1 kg Gender: Female Reproductive Status: Intact Age: Adult

- 1. Cervical vertebra
- Clavicle
 Humerus
- 4. Coracoid
- 5. Scapula
- 6. Rib7. Uncinate process of rib
- 8. Thoracic vertebra
- 9. Synsacrum
- 10. Ilium
- 11. Head of femur 12. (Ilioischiadiac foramen)
- 13. Ischium
- 14. Pubis
- 15. Femur

- 16. (Obturator foramen)
- 17. (Patella)
- 18. Tibiotarsal bone
- 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae
- 22. Pygostyle
- 23. Apex carinae
- 24. Dorsal tubercle of humerus
- 25. Head of humerus
- 26. Ventral tubercle of humerus
- 27. Trochanter of femur




Figure 10-7, A-B
Type of Bird: Blue and Gold Macaw
Type of Study: Wing
Projection: Mediolateral
Weight of Bird: 1.1 kg Gender: Female

Reproductive Status: Intact Age: Adult

- $1. \ Head \ of \ humerus$
- 2. Dorsal tubercle of humerus
- 3. Dorsal condyle of humerus
- 4. Radius
- 5. Radial carpal bone
- 6. Alula7. Major metacarpal bone
- 8. Phalanges of major digit
- 9. Ventral tubercle of humerus
- 10. Humerus
- 11. Ventral condyle of humerus
- 12. Ulna
- 13. Ulnar carpal bone
- 14. Minor metacarpal bone
- 15. Phalanges of minor digit




Figure 10-8, A-BType of Bird: Blue and Gold Macaw Type of Study: Wing Projection: Caudocranial Weight of Bird: 1.1 kg Gender: Female Reproductive Status: Intact Age: Adult

- 1. Head of humerus
- 2. Ventral tubercle of humerus3. Ventral condyle of humerus
- 4. Radius
- 5. Radial carpal bone6. Alula
- 7. Minor metacarpal bone
- 8. Phalanges of major digit
- 9. Dorsal tubercle of humerus
- 10. Humerus
- 11. Dorsal condyle of humerus
- 12. Ulna
- 13. Ulnar carpal bone
- 14. Major metacarpal bone15. Phalanges of minor digit




Figure 10-9, A-BType of Bird: Blue and Gold Macaw Type of Study: Pelvic limb Projection: Mediolateral Weight of Bird: 1.1 kg Gender: Female Reproductive Status: Intact Age: Adult

- Femur
 Patella
- 3. Condyles of femur
- 4. Fibula
- 5. Tibiotarsal bone
- 6. Condyles of tibiotarsal bone
- 7. (Hypotarsal crest of tarsometatarsal bone)
- 8. (Tarsometatarsal bone)
- 9. (Trochlea of tarsometatarsal bone) 10. (Digit I)
- 11. (Digit II)

- 12. (Digit III) 13. (Digit IV) 14. Phalanges 15. (Metatarsal I)




Figure 10-10, A-B Type of Bird: Blue and Gold Macaw Type of Study: Pelvic limb Projection: Craniocaudal Weight of Bird: 1.1 kg Gender: Female Reproductive Status: Intact Age: Adult

- 1. Trochanter of femur
- 2. Femur
- 3. Lateral condyle of femur
- 4. Fibula
- 5. Lateral condyle of tibiotarsal bone
- 6. Tarsometatarsal bone
- 7. Trochlea of tarsometatarsal bone
- 8. Digit IV 9. Digit III
- 10. Head of femur
- 11. Medial condyle of femur

- 12. Tibiotarsal bone
- 13. Medial condyle of tibiotarsal bone
- 14. Digit I 15. Digit II
- 16. Phalanges
- 17. (Patella)
- 18. (Hypotarsal crest of tarsometatarsal bone)
 19. Metatarsal I




Figure 10-11, A-B
Type of Bird: Blue and Gold Macaw Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)

30 ml administered via gavage tube Projection: Laterolateral (right lateral

recumbency)
Weight of Bird: 1.1 kg Gender: Female Reproductive Status: Intact Age: Adult

lmage	Time (hr)
A	Scout
В	Immediate

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 10-11, C-D

Type of Bird: Blue and Gold Macaw Type of Study: Gastrointestinal positive contrast study

Contrast Study
Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 30 ml administered via gavage tube
Projection: Laterolateral (right lateral

recumbency)
Weight of Bird: 1.1 kg

Gender: Female Reproductive Status: Intact Age: Adult

lmage	Time (hr)
С	0.17
D	1.0

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. Large intestine
- 8. Cloaca
- 9. Proventricular-ventricular isthmus




Figure 10-11, E

Type of Bird: Blue and Gold Macaw

Type of Study: Gastrointestinal positive contrast study
Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutica, Inc., Lafayette, IN) 30 ml administered via gavage tube
Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 1.1 kg

Gender: Female

Reproductive Status: Intact Age: Adult

Image	Time (hr)
E	2.0

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
 5. Intestines
- 6. Duodenum
- 7. Large intestine
- 8. Cloaca




Figure 10-12, A-B

Type of Bird: Blue and Gold Macaw
Type of Study: Gastrointestinal positive
contrast study

Contrast Study
Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 30 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 1.1 kg Gender: Female Reproductive Status: Intact Age: Adult

Image	Time (hr)
A	Scout
В	Immediate

- 1. Crop
- 2. Esophagus
- 3. Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
- 8. (Cloaca)
- 9. (Proventricular-ventricular isthmus)




Figure 10-12, C-DType of Bird: Blue and Gold Macaw
Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 30 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 1.1 kg Gender: Female Reproductive Status: Intact Age: Adult

lmage	Time (hr)
С	0.17
D	1.0

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)
- 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 10-12, E
Type of Bird: Blue and Gold Macaw
Type of Study: Gastrointestinal positive
contrast study Contrast Medium: Barium sulfate suspen-

sion (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 30 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 1.1 kg Gender: Female Reproductive Status: Intact Age: Adult

lmage	Time (hr)
Е	2.0

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. Large intestine
- 8. Cloaca
- 9. Proventricular-ventricular isthmus




Figure 10-13, A-D Type of Bird: Blue and Gold Macaw Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.3 kg Age: Adult

- 2. Mandible 3. Hyoid bone 4. (Endotracheal tube) 5. (Keratinized maxillary beak) 6. (Nasal cavity) 7. Nasal concha 8. (Pharynx) 9. (Premaxillary bone)
 10. (Infraorbital sinus)
- 11. (Nasal septum) 12. (Choana) 13. Palatine bone 14. (Sphenoid bone)
- 15. (Jugal [zygomatic] bone) 16. (Frontal bone)
- 17. (Pterygoid bone) 18. (Eyeball)

- 19. (Scleral ossicle)20. (Interorbital septum)
- 21. (Lens of eyeball)
- 22. (Trachea)
 23. (Cerebrum)
- 24. (External ear canal)
- 25. (Cerebellum)
- 26. (Spinal cord) 27. (Dens)
- 28. (Cere)
- 29. (Nare[s])
- 30. (Feather)
- 31. (Pons)
 32. (Occipital bone)
- 33. (Cervical vertebra)




Figure 10-13, E-H

Type of Bird: Blue and Gold Macaw Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.3 kg Age: Adult

- 1. (Tongue)
- 2. Mandible
- 3. Hyoid bone
- 4. (Endotracheal tube)
- 5. (Keratinized maxillary beak)
- 6. Nasal cavity
- 7. (Nasal concha)
- 8. Pharynx
- 9. (Premaxillary bone)
 10. (Infraorbital sinus)
- 11. (Nasal septum)
- 12. Choana
- 13. Palatine bone
- 14. (Sphenoid bone)
- 15. (Jugal [zygomatic] bone)
- 16. (Frontal bone)
- 17. (Pterygoid bone)
- 18. Eyeball

- 19. (Scleral ossicle)
- 20. (Interorbital septum)
- 21. (Lens of eyeball)
- 22. (Trachea)
- 23. (Cerebrum)
- 24. (External ear canal)
- 25. (Cerebellum)
- 26. (Spinal cord) 27. (Dens)
- 28. (Cere)
- 29. (Nare[s])
- 30. (Feather)
- 31. (Pons)
 32. (Occipital bone)
- 33. (Cervical vertebra)




Figure 10-13, I-L Type of Bird: Blue and Gold Macaw Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.3 kg Age: Adult

- 1. (Tongue) 2. Mandible 3. (Hyoid bone)
- 4. (Endotracheal tube)
 5. (Keratinized maxillary beak)
- 6. (Nasal cavity) 7. (Nasal co 8. Pharynx (Nasal concha)
- 9. (Premaxillary bone) 10. Infraorbital sinus
- 11. (Nasal septum)
- 12. (Choana) 13. Palatine bone
- 14. (Sphenoid bone)
- 15. Jugal [zygomatic] bone
- 16. (Frontal bone)
- 17. (Pterygoid bone)
- 18. Eyeball

- 19. Scleral ossicle
- 20. (Interorbital septum)
- 21. (Lens of eyeball)
- 22. (Trachea)
- 23. Cerebrum
- 24. (External ear canal)
- 25. (Cerebellum)
- 26. (Spinal cord) 27. (Dens)
- 28. (Cere)
- 29. (Nare[s])
- 30. (Feather)
- 31. (Pons) 32. (Occipital bone)
- 33. (Cervical vertebra)

NOTE: Structures in parentheses are not labeled.





Figure 10-13, M-P

Type of Bird: Blue and Gold Macaw Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.3 kg Age: Adult

- 1. (Tongue)
- 2. Mandible
- 3. (Hyoid bone)
- 4. Endotracheal tube
- 5. (Keratinized maxillary beak)
- 6. (Nasal cavity)
- 7. (Nasal concha)
- 8. (Pharynx)
- 9. (Premaxillary bone)
- 10. (Infraorbital sinus)
- 11. (Nasal septum)
- 12. (Choana)
- 13. (Palatine bone)
- 14. (Sphenoid bone)
- 15. (Jugal [zygomatic] bone)
- 16. (Frontal bone)
- 17. (Pterygoid bone)
- 18. (Eyeball)

- 19. (Scleral ossicle)
- 20. (Interorbital septum)
- 21. (Lens of eyeball)
- 22. Trachea
- 23. Cerebrum
- 24. (External ear canal)
- 25. Čerebellum
- 26. Spinal cord
- 27. (Dens)
- 28. (Cere)
- 29. (Nare[s])
- 30. (Feather)
- 31. (Pons)
- 32. (Occipital bone) 33. Cervical vertebra

Goffin Cockatoo (Cacatua goffiniana)






Figure 11-1, A-B

Type of Bird: Goffin Cockatoo Type of Study: Viscera of the coelom
Projection: Laterolateral (right lateral
recumbency)
Weight of Bird: 225 g

Gender: Female

Reproductive Status: Intact

Age: Adult

- 1. Trachea
- 2. (Crop)
- 3. Brachiocephalic artery and aorta
- 4. (Brachiocephalic artery)
- 5. Àorta
- 6. Pulmonary artery
- 7. Pulmonary vein
- 8. Heart
- 9. Left atrium
- 10. Esophagus
- ll. Liver
- 12. Lung
- 13. Syrinx
- 14. (Gonad) 15. Ovary 16. (Testes)

- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. Cloaca
- 21. (Cervical air sac)22. (Clavicular air sac)
- 23. Thoracic air sac
- 24. Abdominal air sac
- 25. (Apex of heart)
- 26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys
- 28. Spleen





Figure 11-2, A-B
Type of Bird: Goffin Cockatoo
Type of Study: Viscera of the coelom
Projection: Ventrodorsal
Weight of Bird: 225 g Gender: Female Reproductive Status: Intact Age: Adult

- Trachea
 Cervical air sac
 Crop
- 4. Clavicular air sac
- 5. (Brachiocephalic artery and aorta)6. Heart base vessel
- 7. Heart
- 8. Lung
- 9. Thoracic air sac

- 10. Liver 11. Kidneys
- 12. (Proventriculus)13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. Cloaca

Sulphur-Crested Cockatoo (Cacatua galerita)






Figure 12-1, A-D Type of Bird: Sulphur-Crested Cockatoo Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 395 g Age: Adult

- Tongue
 Mandible
 Hyoid bone
- 4. Endotracheal tube
- 5. (Keratinized maxillary beak)
- 6. Nasal cavity
- 7. (Nasal concha) 8. Pharynx
- 9. (Premaxillary bone)
- 10. Infraorbital sinus
- 11. Nasal septum
- 12. Choana
- 13. Palatine bone
- 14. (Sphenoid bone)
- 15. (Jugal [zygomatic] bone)
- 16. Frontal bone
- 17. (Pterygoid bone)
 18. (Eyeball)

- 19. Scleral ossicle
- 20. (Interorbital septum)
 21. Lens of eyeball
- 22. (Trachea)
- 23. (Cerebrum)
- 24. (External ear canal)
- 25. (Cerebellum)
- 26. (Spinal cord)
- 27. (Dens)
- 28. Čere
- 29. (Nare[s])
- 30. (Feather)
- 31. (Pons)
- 32. (Occipital bone)
- 33. (Cervical vertebra)

NOTE: Structures in parentheses are not labeled.





Figure 12-1, E-H Type of Bird: Sulphur-Crested Cockatoo Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 395 g Age: Adult

- (Tongue)
 Mandible
 Hyoid bone
- 4. (Endotracheal tube)
- 5. (Keratinized maxillary beak)
- 6. (Nasal cavity)
- (Nasal concha)
- 7. (Nasal co 8. Pharynx
- 9. (Premaxillary bone)
- 10. Infraorbital sinus
- 11. (Nasal septum)
 12. (Choana)
- 13. (Palatine bone)
- 14. (Sphenoid bone)
- 15. Jugal [zygomatic] bone 16. (Frontal bone)
- 17. Pterygoid bone
- 18. Eyeball

- 19. Scleral ossicle
- 20. (Interorbital septum)
- 21. (Lens of eyeball)
- 22. (Trachea)
- 23. Cerebrum
- 24. (External ear canal)
 25. Cerebellum
- 26. (Spinal cord)
- 27. (Dens)
- 28. (Cere) 29. (Nare[s])
- 30. (Feather)
- 31. (Pons)
- 32. (Occipital bone) 33. (Cervical vertebra)

Umbrella Cockatoo (Cacatua alba)






Figure 13-1, A-B

Type of Bird: Umbrella Cockatoo Type of Study: Viscera of the coelom Projection: Laterolateral (right lateral recumbency)

Weight of Bird: 450 g Gender: Female

Reproductive Status: Intact

Age: 3 years

- 1. Trachea
- 2. (Crop)3. Brachiocephalic artery and aorta
- 4. (Brachiocephalic artery)
- 5. Aorta
- 6. Pulmonary artery7. Pulmonary vein
- 8. Heart
- 9. Left atrium
- 10. Esophagus
- 11. Liver
- 12. Lung
- 13. Syrinx
- 14. (Gonad) 15. (Ovary) 16. (Testes)

- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. Cloaca
- 21. Cervical air sac 22. (Clavicular air sac)
- 23. Thoracic air sac 24. (Abdominal air sac)
- 25. (Apex of heart)
- 26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys 28. Spleen





Figure 13-2, A-BType of Bird: Umbrella Cockatoo
Type of Study: Viscera of the coelom Projection: Ventrodorsal Weight of Bird: 450 g Gender: Female Reproductive Status: Intact Age: 3 years

- Trachea
 (Cervical air sac)
 Crop
- 4. Clavicular air sac
- 5. Brachiocephalic artery and aorta6. Heart base vessel
- 7. Heart
- 8. Lung
- 9. Thoracic air sac

- 10. Liver 11. Kidneys
- 12. Proventriculus
- 13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. Cloaca

Moluccan Cockatoo (Cacatua moluccensis)






Figure 14-1, A-B

Type of Bird: Moluccan cockatoo Prandial Condition: Fasted Type of Study: Viscera of the coelom Projection: Laterolateral (right lateral recumbency) Weight of Bird: 800 g Gender: Female

Reproductive Status: Intact

Age: Adult

- 1. Trachea
- 2. (Crop)
- 3. Brachiocephalic artery and aorta
- 4. (Brachiocephalic artery)
- 5. Aorta
- 6. Pulmonary artery
- 7. Pulmonary vein
- 8. Heart
- 9. Left atrium
- 10. Esophagus
- 11. Liver
- 12. Lung
- 13. Syrinx
- 14. (Gonad) 15. Ovary
- 16. (Testes)

- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. Cloaca
- 21. Cervical air sac
- 22. (Clavicular air sac)
- 23. Thoracic air sac
- 24. (Abdominal air sac)
- 25. (Apex of heart)
- 26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys
- 28. Spleen




Figure 14-2, A-BType of Bird: Moluccan Cockatoo Prandial Condition: Fasted Type of Study: Viscera of the coelom Projection: Ventrodorsal Weight of Bird: 800 g Gender: Female Reproductive Status: Intact Age: Adult

- 1. Trachea

- (Cervical air sac)
 (Crop)
 Clavicular air sac
 (Brachiocephalic artery and aorta)
 (Heart base vessel)
- 7. Heart
- 8. Lung
 9. Thoracic air sac
 10. Liver

- 11. Kidneys12. Proventriculus
- 13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. Cloaca




Figure 14-3, A-B

Type of Bird: Moluccan Cockatoo Prandial Condition: Fed (pelleted diet) Type of Study: Viscera of the coelom Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 800 g

Gender: Female

Reproductive Status: Intact

Age: Adult

- 1. Trachea
- 2. Crop3. Brachiocephalic artery and aorta
- 4. (Brachiocephalic artery)
- 5. Aorta
- 6. Pulmonary artery
 7. Pulmonary vein
 8. Heart

- 9. Left atrium
- 10. Esophagus
- 11. Liver
- 12. Lung
- 13. Syrinx
- 14. (Gonad) 15. Ovary 16. (Testes)

- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. Cloaca
- 21. Cervical air sac
- 22. (Clavicular air sac)
- 23. Thoracic air sac
- 24. (Abdominal air sac)
- 25. Apex of heart
- 26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys
- 28. (Spleen)




Figure 14-4, A-B
Type of Bird: Moluccan Cockatoo
Prandial condition: Fed (pelleted diet) Type of Study: Viscera of the coelom Projection: Ventrodorsal Weight of Bird: 800 g Gender: Female Reproductive Status: Intact Age: Adult

- Trachea
 (Cervical air sac)
 (Crop)
- 4. Clavicular air sac
- 5. Brachiocephalic artery and aorta6. (Heart base vessel)7. Heart

- 8. Lung
- 9. Thoracic air sac
- 10. Liver

- 11. Kidneys12. Proventriculus
- 13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. Cloaca




Figure 14-5, A-BType of Bird: Moluccan Cockatoo

Type of Study: Head
Projection: Laterolateral (right lateral recumbency)

Weight of Bird: 800 g Gender: Female

Reproductive Status: Intact Age: Adult

- 1. Keratinized maxillary beak
- 2. Keratinized mandibular beak
- 3. Premaxillary bone
- 4. Mandible
- 5. Hyoid bones
- 6. Palatine bone
- 7. (Pterygoid bone)
 8. External nares
 9. Jugal [zygomatic] bone
- 10. Craniofacial flexion zone

- 11. Orbit
- 12. Frontal bone
- 13. Cranium
- 14. Temporal bone
- 15. Quadrate bone
- 16. Parietal bone




Figure 14-6, A-B
Type of Bird: Moluccan Cockatoo
Type of Study: Head
Projection: Ventrodorsal
Weight of Bird: 800 g
Gender: Female
Reproductive Status: Intact
Age: Adult

- Keratinized maxillary beak
 Premaxillary bone
 Mandible
 Orbit

- 5. Jugal [zygomatic] bone6. Scleral ossicles7. Quadrate bone

- 8. Cranium




Figure 14-7, A-B

Age: Adult

Type of Bird: Moluccan Cockatoo Type of Study: Whole body skeleton Projection: Laterolateral (right lateral recumbency) Weight of Bird: 800 g Gender: Female Reproductive Status: Intact

- 1. Cervical vertebra
- 2. Clavicle
- 3. Humerus 4. Coracoid
- 5. Scapula
- 6. Rib
- 7. (Uncinate process of rib)
 8. Thoracic vertebra
- 9. Synsacrum
- 10. Ilium
- 11. Head of femur
- 12. Ilioischiadiac foramen
- 13. Ischium

- 14. Pubis
- 15. Femur
- 16. (Obturator foramen)
- 17. Patella
- 18. Tibiotarsal bone
- 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae
- 22. Pygostyle




Figure 14-8, A-BType of Bird: Moluccan Cockatoo Type of Study: Whole body skeleton Projection: Ventrodorsal Weight of Bird: 800 g Gender: Female Reproductive Status: Intact Age: Adult

Α

- 1. Cervical vertebra
- 2. Clavicle
- 3. Humerus
- 4. Coracoid 5. Scapula 6. Rib

- 7. (Uncinate process of rib)8. Thoracic vertebra
- 9. Synsacrum
- 10. Ilium
 11. Head of femur
- 12. (Ilioischiadiac foramen)
- 13. İschium
- 14. Pubis
- 15. Femur

- 16. (Obturator foramen)
- 17. (Patella)
- 18. Tibiotarsal bone
- 19. Fibula
- 20. Sternum
 21. Caudal vertebrae
- 22. Pygostyle
- 23. Apex carinae
- 24. Dorsal tubercle of humerus
- 25. Head of humerus26. Ventral tubercle of humerus
- 27. Trochanter of femur




Figure 14-9, A-B Type of Bird: Moluccan Cockatoo
Type of Study: Wing
Projection: Mediolateral
Weight of Bird: 800 g

Gender: Female

Reproductive Status: Intact Age: Adult

1. Head of humerus

- 2. Dorsal tubercle of humerus
- 3. Dorsal condyle of humerus
- 4. Radius
- 5. Radial carpal bone
- 6. Alula7. Major metacarpal bone
- 8. Phalanges of major digit
- 9. Ventral tubercle of humerus
- 10. Humerus
- 11. Ventral condyle of humerus
- 12. Ulna
- 13. Ulnar carpal bone
- 14. Minor metacarpal bone
- 15. Phalanges of minor digit




Figure 14-10, A-B
Type of Bird: Moluccan Cockatoo
Type of Study: Wing Projection: Caudocranial Weight of Bird: 800 g Gender: Female Reproductive Status: Intact Age: Adult

- 1. Head of humerus
- 2. Ventral tubercle of humerus
- 3. Ventral condyle of humerus
- 4. Radius
- 5. Radial carpal bone
- 6. Alula 7. Minor metacarpal bone
- 8. Phalanges of major digit
- 9. Dorsal tubercle of humerus
- 10. Humerus
- 11. Dorsal condyle of humerus 12. Ulna
- 13. (Ulnar carpal bone)
- 14. Major metacarpal bone 15. Phalanges of minor digit




Figure 14-11, A-B

Type of Bird: Moluccan Cockatoo Type of Study: Pelvic limb Projection: Mediolateral Weight of Bird: 800 g Gender: Female Reproductive Status: Intact Age: Adult

- 1. Femur
- 2. Patella 3. Condyles of femur4. Fibula5. Tibiotarsal bone

- 6. Condyles of tibiotarsal bone
- 7. Hypotarsal crest of tarsometatarsal bone
- 8. Tarsometatarsal bone
- 9. Trochlea of tarsometatarsal bone

- 10. Digit I 11. Digit II 12. Digit III
- 13. Digit IV
- 14. Phalanges15. Metatarsal I




Figure 14-12, A-B Type of Bird: Moluccan Cockatoo Type of Study: Pelvic limb Projection: Craniocaudal Weight of Bird: 800 g Gender: Female Reproductive Status: Intact Age: Adult

- 1. Trochanter of femur
- 2. Femur
- 3. Lateral condyle of femur
- 4. Fibula
- 5. Lateral condyle of tibiotarsal bone
- 6. Tarsometatarsal bone
- 7. Trochlea of tarsometatarsal bone
- 8. Digit IV 9. Digit III
- 10. Head of femur
- 11. Medial condyle of femur

- 12. Tibiotarsal bone
- 13. Medial condyle of tibiotarsal bone
- 14. Digit I
- 15. Digit II
- 16. Phalanges
- 17. (Patella)
- 18. Hypotarsal crest of tarsometatarsal bone
- 19. Metatarsal I




Figure 14-13, A-B Type of Bird: Moluccan Cockatoo
Type of Study: Distal pelvic limb
Projection: Mediolateral
Weight of Bird: 800 g
Gender: Female
Reproductive Status: Intact

Age: Adult

- 1. Tarsometatarsal bone
- 2. Trochlea of tarsometatarsal bone 3. Metatarsal bone I

- 4. Digit I
 5. Digit II
 6. Digit III

- 7. Digit IV
- 8. Phalanges
- 9. Hypotarsal crest of tarsometatarsal bone
 10. Condyles of tibiotarsal bone




Figure 14-14, A-B
Type of Bird: Moluccan Cockatoo
Type of Study: Distal pelvic limb
Projection: Dorsoplantar
Weight of Bird: 800 g
Gender: Female Reproductive Status: Intact Age: Adult

- 1. Tarsometatarsal bone
- 2. Trochlea of tarsometatarsal bone
- Hochiea of carson
 Metatarsal bone I
 Digit I
 Digit III
 Digit III

- 7. Digit IV
- 8. Phalanges
 9. Hypotarsal crest of tarsometatarsal bone
- 10. Condyles of tibiotarsal bone




Figure 14-15, A-B

Type of Bird: Moluccan Cockatoo Type of Study: Gastrointestinal positive contrast study

Contrast Study:
Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 25 ml administered via gavage tube Projection: Laterolateral (right lateral recumbency)

Weight of Bird: 820 g Gender: Female Reproductive Status: Intact Age: Adult

Time (hr) Image Α Scout В 0.25

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 14-16, A-B

Type of Bird: Moluccan Cockatoo
Type of Study: Gastrointestinal positive
contrast study

contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)

25 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 820 g Gender: Female

Reproductive Status: Intact

Age: Adult

Image	Time (hr)
A	Scout
В	0.25

- 1. Crop
- 2. Esophagus
- 3. Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
- 8. Cloaca
- 9. Proventricular-ventricular isthmus




Figure 14-17, A-F Type of Bird: Moluccan Cockatoo Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 820 g Age: Adult

- 2. Mandible 3. Hyoid bone 4. (Endotracheal tube) 5. Keratinized maxillary beak 6. (Nasal cavity)7. Nasal concha 8. Pharynx 9. (Premaxillary bone) 10. Infraorbital sinus 11. Nasal septum
- 12. Choana 13. Palatine bone 14. (Sphenoid bone) 15. (Jugal [zygomatic] bone) 16. (Frontal bone)
- 17. (Pterygoid bone)
 18. (Eyeball)

1. Tongue

- 19. (Scleral ossicle)
- 20. (Interorbital septum)
 21. (Lens of eyeball)
- 22. (Trachea)
- 23. (Cerebrum)
- 24. (External ear canal)
- 25. (Cerebellum) 26. (Spinal cord)
- 27. (Dens)
- 28. Cere
- 29. Nare[s]
- 30. (Feather)
- 31. (Pons) 32. (Occipital bone)
- 33. (Cervical vertebra)




Figure 14-17, G-L Type of Bird: Moluccan Cockatoo Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 820 g Age: Adult

- Tongue
 Mandible
 Hyoid bone
- 4. (Endotracheal tube)
- 5. (Keratinized maxillary beak)
- 6. (Nasal cavity)

- 7. (Nasal concha)
 8. (Pharynx)
 9. (Premaxillary bone)
 Conference of the fraction of the fracti
- 11. Nasal septum
- 12. Choana
- 13. Palatine bone
- 14. (Sphenoid bone)
- 15. Jugal [zygomatic] bone 16. Frontal bone
- 17. Pterygoid bone 18. Eyeball

- 19. Scleral ossicle
- 20. Interorbital septum
 21. Lens of eyeball
- 22. (Trachea)
- 23. Cerebrum
- 24. (External ear canal)
- 25. (Cerebellum)
- 26. (Spinal cord)
- 27. (Dens)
- 28. Čere
- 29. (Nare[s]) 30. Feather
- 31. (Pons)
- 32. (Occipital bone)
 33. (Cervical vertebra)




Figure 14-17, M-OType of Bird: Moluccan Cockatoo

Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 820 g Age: Adult

- (Tongue)
 (Mandible) 3. (Hyoid bone)
 4. (Endotracheal tube)
 5. (Keratinized maxillary beak)
- 6. (Nasal cavity)
- (Nasal concha) 8. (Pharynx)
- 9. (Premaxillary bone) 10. (Infraorbital sinus)
- 11. (Nasal septum)
- 12. (Choana)
- 13. (Palatine bone)
 14. (Sphenoid bone)
- 15. (Jugal [zygomatic] bone)
- 16. (Frontal bone)
- 17. (Pterygoid bone)
 18. (Eyeball)

- 19. (Scleral ossicle)
- 20. (Interorbital septum)
 21. (Lens of eyeball)

- 22. (Trachea)
 23. (Cerebrum)
- 24. (External ear canal)
- 25. Cerebellum
- 26. (Spinal cord)
 27. (Dens)
 28. (Cere)

- 29. (Nare[s])
- 30. Feather
- 31. Pons
- 32. Occipital bone
- 33. Cervical vertebra

Pigeon (Columba livia)






Figure 15-1, A-B

Type of Bird: Pigeon Type of Study: Viscera of the coelom Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 545 g

Gender: Unknown

Reproductive Status: Intact

Age: Adult

- 1. Trachea
- 2. Crop
- 3. Brachiocephalic artery and aorta
- 4. (Brachiocephalic artery)
- 5. Aorta
- 6. Pulmonary artery
- 7. Pulmonary vein
- 8. Heart
- 9. Left atrium
- 10. Esophagus
- 11. Liver
- 12. Lung
- 13. Syrinx
- 14. Gonad
- 15. (Ovary)

- 16. (Testes)
- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. Cloaca
- 21. Cervical air sac
- 22. (Clavicular air sac)
- 23. Thoracic air sac
- 24. Abdominal air sac 25. (Apex of heart)
- 26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys
- 28. Spleen




Figure 15-2, A-B Type of Bird: Pigeon Type of Study: Viscera of the coelom Projection: Ventrodorsal Weight of Bird: 545 g Gender: Unknown Reproductive Status: Intact Age: Adult

- 1. Trachea
- 2. (Cervical air sac)

- 3. Crop4. Clavicular air sac5. Brachiocephalic artery and aorta
- 6. Heart base vessel
- 7. Heart
- 8. Lung
 9. Thoracic air sac

- 10. Liver 11. Kidneys
- 12. Proventriculus
- 13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. Cloaca




Figure 15-3, A-B

Age: Adult

Type of Bird: Pigeon
Type of Study: Head
Projection: Laterolateral (right lateral recumbency) Weight of Bird: 590 g Gender: Unknown Reproductive Status: Intact

- 1. Keratinized maxillary beak
- Keratinized mandibular beak
 Fremaxillary bone
 Mandible

- 5. Hyoid bones
- 6. Palatine bone
- 7. (Pterygoid bone)
- 8. External nares
- 9. Jugal [zygomatic] bone
- 10. Craniofacial flexion zone

- 11. Orbit
- 12. Frontal bone
- 13. Cranium
- 14. Temporal bone
- 15. Quadrate bone
- 16. Parietal bone




Figure 15-4, A-B
Type of Bird: Pigeon
Type of Study: Head
Projection: Ventrodorsal
Weight of Bird: 590 g
Gender: Unknown Reproductive Status: Intact Age: Adult

- Keratinized maxillary beak
 Premaxillary bone
 (Mandible)
 Orbit

- 5. (Jugal [zygomatic] bone)




Figure 15-5, A-BType of Bird: Pigeon Type of Study: Whole body skeleton Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 590 g
Gender: Unknown

Reproductive Status: Intact Age: Adult

- 1. Cervical vertebra
- 2. Clavicle
- 3. Humerus 4. Coracoid
- 5. Scapula
- 6. Rib
- 7. (Uncinate process of rib)
- 8. Thoracic vertebra
- 9. Synsacrum
- 10. (Ilium) 11. Head of femur
- 12. (Ilioischiadiac foramen)

- 13. (Ischium) 14. Pubis
- 15. Femur
- 16. (Obturator foramen)
- 17. Patella18. Tibiotarsal bone 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae
- 22. Pygostyle




Figure 15-6, A-B Type of Bird: Pigeon Type of Study: Whole body skeleton Projection: Ventrodorsal Weight of Bird: 590 g Gender: Unknown Reproductive Status: Intact Age: Adult

- 1. Cervical vertebra
- 2. Clavicle
 3. (Humerus)
- 4. Coracoid 5. Scapula
- 6. Rib
- 7. Uncinate process of rib8. Thoracic vertebra
- 9. Synsacrum
- 10. Ilium
- 11. Head of femur
- 12. (Ilioischiadiac foramen)
 13. Ischium
- 14. Pubis
- 15. Femur

- 16. (Obturator foramen)17. (Patella)18. Tibiotarsal bone

- 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae
- 22. Pygostyle
 23. Apex carinae
- 24. Dorsal tubercle of humerus
- 25. Head of humerus
- 26. Ventral tubercle of humerus
- 27. Trochanter of femur




Figure 15-7, A-BType of Bird: Pigeon Type of Study: Wing Projection: Mediolateral Weight of Bird: 590 g Gender: Unknown

Reproductive Status: Intact

Age: Adult

- 2. Dorsal tubercle of humerus
- $3. \ Dorsal\ condyle\ of\ humerus$
- 4. Radius
- 5. Radial carpal bone
- 6. Alula7. Major metacarpal bone
- 8. Phalanges of major digit
- 10. Humerus
- 11. Ventral condyle of humerus
- 12. Ulna

- 13. Ulnar carpal bone14. Minor metacarpal bone15. Phalanges of minor digit




Figure 15-8, A-B
Type of Bird: Pigeon
Type of Study: Wing Projection: Caudocranial Weight of Bird: 590 g Gender: Unknown Reproductive Status: Intact Age: Adult

- Head of humerus
 Ventral tubercle of humerus
 Ventral condyle of humerus
- 4. Radius
- 5. Radial carpal bone
- 6. Alula
- 7. Minor metacarpal bone8. Phalanges of major digit
- 9. Dorsal tubercle of humerus
- 10. Humerus
- 11. Dorsal condyle of humerus
- 12. Ulna

- 13. (Ulnar carpal bone)14. Major metacarpal bone15. Phalanges of minor digit




Figure 15-9, A-B
Type of Bird: Pigeon
Type of Study: Pelvic limb
Projection: Mediolateral
Weight of Bird: 590 g Gender: Unknown Reproductive Status: Intact Age: Adult

- Femur
 Patella
- 3. Condyles of femur4. Fibula
- 5. Tibiotarsal bone
- 6. Condyles of tibiotarsal bone
- 7. Hypotarsal crest of tarsometatarsal bone
- 8. Tarsometatarsal bone
- 9. (Trochlea of tarsometatarsal bone)

- 10. Digit I
 11. Digit II
 12. Digit III
 13. Digit IV
 14. Phalanges
- 15. Metatarsal I




Figure 15-10, A-B Type of Bird: Pigeon
Type of Study: Pelvic limb Projection: Craniocaudal Weight of Bird: 590 g Gender: Unknown Reproductive Status: Intact Age: Adult

- 1. Trochanter of femur
- 2. Femur
- 3. Lateral condyle of femur
- 4. Fibula
- 5. Lateral condyle of tibiotarsal bone
- 6. Tarsometatarsal bone
- 7. Trochlea of tarsometatarsal bone

- 8. Digit IV
 9. Digit III
 10. Head of femur
- 11. Medial condyle of femur

- 12. Tibiotarsal bone
- 13. Medial condyle of tibiotarsal bone

- 14. Digit I 15. Digit II 16. Phalanges
- 17. (Patella)
- 18. (Hypotarsal crest of tarsometatarsal bone)
- 19. Metatarsal I




Figure 15-11, A-B
Type of Bird: Pigeon
Type of Study: Distal pelvic limb
Projection: Mediolateral
Weight of Bird: 590 g

Gender: Unknown Reproductive Status: Intact Age: Adult

- 1. Tarsometatarsal bone
- 2. Trochlea of tarsometatarsal bone
- 3. Metatarsal bone I
- 4. Digit I 5. Digit II

- 6. Digit III 7. Digit IV

- 8. Phalanges
 9. Hypotarsal crest of tarsometatarsal bone
- 10. Condyles of tibiotarsal bone




Figure 15-12, A-B
Type of Bird: Pigeon
Type of Study: Distal pelvic limb
Projection: Dorsoplantar
Weight of Bird: 590 g Gender: Unknown Reproductive Status: Intact Age: Adult

- 1. Tarsometatarsal bone
- 2. Trochlea of tarsometatarsal bone
- Froctilea of tarson
 Metatarsal bone I
 Digit I
 Digit II

- 6. Digit III
 7. Digit IV
 8. Phalanges
 9. (Hypotarsal crest of tarsometatarsal bone)
- 10. Condyles of tibiotarsal bone




Figure 15-13, A-B
Type of Bird: Pigeon
Type of Study: Respiratory series
Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 545 g
Gender: Unknown
Reproductive Status: Intact
Age: Adult

Image	Phase of Respiration
A	Inspiration, spontaneous ventilation
В	Inspiration, manual assistance (5 cm H ₂ O)



Figure 15-13, C
Type of Bird: Pigeon
Type of Study: Respiratory series
Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 545 g
Gender: Unknown
Reproductive Status: Intact
Age: Adult

Image	Phase of Respiration
С	Inspiration, ventilator assisted (8 cm H2O)




Figure 15-14, A-B

Type of Bird: Pigeon
Type of Study: Gastrointestinal positive

contrast study
Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 20 ml administered via gavage tube

Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 545 g Gender: Unknown Reproductive Status: Intact Age: Adult

Image	Time (hr)
A	Scout
В	0.25

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 15-14, C-D
Type of Bird: Pigeon
Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 20 ml administered via gavage tube Projection: Laterolateral (right lateral

recumbency)
Weight of Bird: 545 g
Gender: Unknown Reproductive Status: Intact Age: Adult

Image	Time (hr)
С	1.0
D	2.0

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)
- 8. (Cloaca)
- 9. (Proventricular-ventricular isthmus)




Figure 15-14, E

Type of Bird: Pigeon
Type of Study: Gastrointestinal positive

contrast study
Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)
20 ml administered via gavage tube
Projection: Laterolateral (right lateral

recumbency) Weight of Bird: 545 g Gender: Unknown

Reproductive Status: Intact

Age: Adult

Image	Time (hr)
Е	3.0

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus
 5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
- 8. (Cloaca)
- 9. (Proventricular-ventricular isthmus)




Figure 15-15, A-B

Type of Bird: Pigeon
Type of Study: Gastrointestinal positive
contrast study

Contrast Study
Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette
Pharmaceutical, Inc., Lafayette, IN)
20 ml administered via gavage tube
Projection: Ventrodesval

Projection: Ventrodorsal Weight of Bird: 545 g Gender: Unknown Reproductive Status: Intact

Age: Adult

lmage	Time (hr)
A	Scout
В	0.25

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)
 8. (Cloaca)
- 9. (Proventricular-ventricular isthmus)




Figure 15-15, C-D

Type of Bird: Pigeon
Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 20 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 545 g Gender: Unknown Reproductive Status: Intact

Age: Adult

Image	Time (hr)
С	1.0
D	2.0

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)
 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 15-15, E

Type of Bird: Pigeon
Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 20 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 545 g Gender: Unknown Reproductive Status: Intact Age: Adult

lmage	Time (hr)
Е	3.0

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)
 8. (Cloaca)
- 9. Proventricular-ventricular isthmus




Figure 15-16

Type of Bird: Pigeon
Type of Study: Cloacagram (retrograde, positive contrast)

Contrast Medium: Diatrizoate meglumine, diatrizoate sodium (RenoCal-76® 37% organically bound iodine; Bracco Diagnostics, Inc., Princeton, NJ) 5 ml administered via vent

Projection: Laterolateral (right lateral

recumbency) Weight of Bird: 545 g Gender: Unknown Reproductive Status: Intact

Age: Adult

lmage	Time (hr)
15-16	Immediate

- 1. Colon
- 2. Coprodeum3. Urodeum
- 4. Cloaca 5. Vent
- 6. Intestines




Figure 15-17
Type of Bird: Pigeon
Type of Study: Cloacagram (retrograde, positive contrast)
Contract Medium Distripate magluming

Contrast Medium: Diatrizoate meglumine, diatrizoate sodium (RenoCal–76® 37% organically bound iodine; Bracco Diagnostics, Inc., Princeton, NJ) 5 ml administered via vent

Projection: Ventrodorsal
Weight of Bird: 545 g
Gender: Unknown
Reproductive Status: Intact


Age: Adult

lmage	Time (hr)
15-17	Immediate

- Colon
 Coprodeum
 Urodeum
 Cloaca
 Vent

- 6. Intestines

Mallard Duck (Anas platyrhynchos)






Figure 16-1, A-B

Type of Bird: Mallard Duck Type of Study: Viscera of the coelom
Projection: Laterolateral (right lateral
recumbency)
Weight of Bird: 1.2 kg

Gender: Male

Reproductive Status: Intact

Age: Adult

- 1. Trachea
- 2. Crop
- 3. Brachiocephalic artery and aorta
- 4. (Brachiocephalic artery)
- 5. (Aorta)
- 6. Pulmonary artery
- 7. Pulmonary vein
- 8. Heart
- 9. Left atrium
- 10. (Esophagus)
- 11. Liver
- 12. Lung
- 13. (Syrinx)
 14. (Gonad)
 15. (Ovary)
 16. Testes

- 17. Proventriculus 18. Ventriculus
- 19. Intestines
- 20. (Cloaca) 21. (Cervical air sac)
- 22. (Clavicular air sac)
- 23. Thoracic air sac
- 24. Abdominal air sac
- 25. (Apex of heart)
- 26. (Interface between caudal thoracic and abdominal air sacs)
- 27. Kidneys
- 28. (Spleen)
- 29. Syringeal bulla




Figure 16-2, A-B Type of Bird: Mallard Duck
Type of Study: Viscera of the coelom
Projection: Ventrodorsal
Weight of Bird: 1.2 kg Gender: Male Reproductive Status: Intact Age: Adult

- 1. Trachea
- 1. Hachea
 2. (Cervical air sac)
 3. Crop
 4. Clavicular air sac

- 5. Brachiocephalic artery and aorta
- 6. Heart base vessel
- 7. Heart
- 8. Lung
 9. Thoracic air sac
- 10. Liver

- 11. Kidneys12. (Proventriculus)13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. Cloaca
- 17. Syringeal bulla




Figure 16-3, A-B

Type of Bird: Mallard Duck
Type of Study: Head
Projection: Laterolateral (right lateral recumbency) Weight of Bird: 1.2 kg Gender: Male

Reproductive Status: Intact

Age: Adult

- 1. Keratinized maxillary beak
- 2. (Keratinized mandibular beak)
- 3. Premaxillary bone
- 4. Mandible
- 5. Hyoid bones
- 6. Palatine bone
 7. (Pterygoid bone)
- 8. External nares
- 9. (Jugal [zygomatic] bone)
- 10. Craniofacial flexion zone 11. Orbit
- 12. Frontal bone
- 13. Cranium
- 14. Temporal bone 15. (Quadrate bone)
- 16. Parietal bone



Figure 16-4, A-B
Type of Bird: Mallard Duck
Type of Study: Head
Projection: Ventrodorsal
Weight of Bird: 1.2 kg
Gender: Male
Reproductive Status: Intact
Age: Adult

- 6. (Scleral ossicles)7. (Quadrate bone)8. Cranium




Figure 16-5, A-B

Type of Bird: Mallard Duck
Type of Study: Whole body skeleton
Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 1.2 kg Gender: Male

Reproductive Status: Intact Age: Adult

- 1. Cervical vertebra
- 2. Clavicle
- 3. Humerus 4. Coracoid
- 5. Scapula
- 6. Rib7. Uncinate process of rib
- 8. Thoracic vertebra
 9. Synsacrum
- 10. Ilium
- 11. Head of femur
- 12. Ilioischiadiac foramen
- 13. (Ischium)

- 14. Pubis
- 15. Femur
- 16. (Obturator foramen)
- 17. Patella
- 18. Tibiotarsal bone
- 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae 22. Pygostyle




Figure 16-6, A-B Type of Bird: Mallard Duck
Type of Study: Whole body skeleton
Projection: Ventrodorsal
Weight of Bird: 1.2 kg Gender: Male Reproductive Status: Intact Age: Adult

- 1. Cervical vertebra
- Clavicle
 Humerus
- 4. Coracoid
- 5. Scapula
- 6. Rib
- 7. (Uncinate process of rib)8. Thoracic vertebra
- 9. Synsacrum
- 10. Ilium
- 11. Head of femur
- 12. (Ilioischiadiac foramen)
 13. Ischium
- 14. Pubis
- 15. Femur

- 16. (Obturator foramen)
- 17. (Patella)
 18. Tibiotarsal bone
- 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae
- 22. Pygostyle
- 23. Apex carinae
- 24. (Dorsal tubercle of humerus)
- 25. Head of humerus
- 26. Ventral tubercle of humerus
- 27. Trochanter of femur





Figure 16-7, A-B
Type of Brid: Mallard Duck
Type of Study: Wing
Projection: Mediolateral
Weight of Bird: 1.2 kg Gender: Male Reproductive Status: Intact

Age: Adult

- 1. Head of humerus
- 2. Dorsal tubercle of humerus
- 3. Dorsal condyle of humerus
- 4. Radius
- 5. Radial carpal bone

- 6. Alula
 7. Major metacarpal bone
 8. Phalanges of major digit
- 9. Ventral tubercle of humerus
- 10. Humerus
- 11. Ventral condyle of humerus 12. Ulna
- 13. Ulnar carpal bone
- 14. Minor metacarpal bone 15. Phalanges of minor digit






Figure 16-9, A-B
Type of Bird: Mallard Duck
Type of Study: Pelvic limb Projection: Mediolateral Weight of Bird: 1.2 kg Gender: Male Reproductive Status: Intact Age: Adult

- Femur
 Patella
 Condyles of femur
- 4. Fibula
- 5. Tibiotarsal bone
- 6. Condyles of tibiotarsal bone
 7. Hypotarsal crest of tarsometatarsal
- 8. Tarsometatarsal bone
- 9. Trochlea of tarsometatarsal bone
- 10. (Digit I)
 11. (Digit II)
 12. (Digit III)

- 13. (Digit IV) 14. Phalanges
- 15. Metatarsal I




Figure 16-10, A-B Type of Bird: Mallard Duck Type of Study: Pelvic limb Projection: Craniocaudal Weight of Bird: 1.2 kg Gender: Male Reproductive Status: Intact Age: Adult

- 1. Trochanter of femur
- 2. Femur
- 3. Lateral condyle of femur
- 4. Fibula
- 5. Lateral condyle of tibiotarsal bone6. Tarsometatarsal bone7. Trochlea of tarsometatarsal bone

- 8. Digit IV
 9. Digit III
 10. Head of femur
- 11. Medial condyle of femur

- 12. Tibiotarsal bone
- 13. Medial condyle of tibiotarsal bone
- 14. Digit I
- 15. Digit II 16. Phalanges
- 17. (Patella)
- 18. (Hypotarsal crest of tarsometatarsal bone)
- 19. Metatarsal I




Figure 16-11, A-B
Type of Bird: Mallard Duck
Type of Study: Distal pelvic limb
Projection: Mediolateral Weight of Bird: 1.2 kg Gender: Male Reproductive Status: Intact Age: Adult

- 1. Tarsometatarsal bone
- 2. Trochlea of tarsometatarsal bone
- 3. Metatarsal bone I

- 4. Digit I
 5. Digit II
 6. Digit III
 7. (Digit IV)

- 8. Phalanges
- 9. Hypotarsal crest of tarsometatarsal bone
- 10. Condyles of tibiotarsal bone




Figure 16-12, A-B
Type of Bird: Mallard Duck
Type of Study: Distal pelvic limb
Projection: Dorsoplantar
Weight of Bird: 1.2 kg

Gender: Male Reproductive Status: Intact Age: Adult

- Tarsometatarsal bone
 Trochlea of tarsometatarsal bone
 Metatarsal bone I

- 4. Digit I
 5. Digit II
 6. Digit III
 7. Digit IV

- 8. Phalanges 9. (Hypotarsal crest of tarsometatarsal bone)
 10. (Condyles of tibiotarsal bone)




Figure 16-13, A-C

Type of Bird: Mallard Duck Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 35 ml administered via gavage tube Projection: Laterolateral (right lateral

recumbency) Weight of Bird: 1.2 kg Gender: Male

Reproductive Status: Intact

Age: Adult

Image	Time (hr)
A	Scout
В	0.25
С	1.0

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
 7. Large intestine
 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 16-14, A-B Type of Bird: Mallard Duck

Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 35 ml administered via gavage tube Projection: Ventrodorsal Weight of Bird: 1.2 kg

Gender: Male

Reproductive Status: Intact

Age: Adult

Image	Time (hr)
A	Scout
В	0.25

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. Large intestine
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 16-14, C Type of Bird: Mallard Duck

Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 35 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 1.2 kg Gender: Male

Reproductive Status: Intact

Age: Adult

Image Time (hr) С 1.0

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. Large intestine
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)



Figure 16-15
Type of Bird: Mallard Duck
Type of Study: Excretory urogramcutaneous ulnar vein contrast medium administration

administration
Contrast Medium: Diatrizoate meglumine, diatrizoate sodium (RenoCal-76® 37% organically bound iodine; Bracco Diagnostics, Inc., Princeton, NJ) 1.2 ml IV (1 ml/kg) via cutaneous ulnar vein

Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 1.2 kg

Gender: Male Reproductive Status: Intact Age: Adult

lmage	Time (min)
16-15	Immediate

- Kidneys
 Ureter
 Cloaca




Figure 16-16
Type of Bird: Mallard Duck
Type of Study: Excretory urogram-medial metatarsal vein contrast administration.
Contrast Medium: Diatrizoate meglumine, diatrizoate sodium (RenoCal-76® 37% organically bound iodine; Bracco Diagnostics, Inc., Princeton, NJ) 1.2 ml IV (1 ml/kg) via medial metatarsal vein Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 1.2 kg

Gender: Male
Reproductive Status: Intact
Age: Adult

lmage	Time (min)
16-16	Immediate

Kidneys
 Ureter
 Cloaca




Figure 16-17, A-D

Type of Bird: Mallard Duck Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.4 kg Age: Adult

- 1. Tongue
- 2. Mandible
- 3. Hyoid bone
- 4. Endotracheal tube
- 5. Keratinized maxillary beak
- 6. (Nasal cavity)
- 7. Nasal concha
- 8. (Pharynx)
- 9. Premaxillary bone
- 10. (Infraorbital sinus)
- 11. (Nasal septum)
- 12. (Choana)
- 13. Palatine bone
 14. (Sphenoid bone)
- 15. Jugal [zygomatic] bone
- 16. (Frontal bone)
- 17. (Pterygoid bone)
 18. (Eyeball)

- 19. (Scleral ossicle)
- 20. (Interorbital septum)
- 21. (Lens of eyeball)
- 22. (Trachea)
- 23. (Cerebrum)
- 24. (External ear canal)
- 25. (Cerebellum)
- 26. (Spinal cord) 27. (Dens)
- 28. (Cere)
- 29. (Nare[s])
- 30. (Feather)
- 31. (Pons)
- 32. (Occipital bone) 33. (Cervical vertebra)




Figure 16-17, E-H

Type of Bird: Mallard Duck Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.4 kg Age: Adult

- 1. Tongue
- 2. Mandible
- 3. Hyoid bone
- 4. (Endotracheal tube)
- 5. (Keratinized maxillary beak)
- 6. (Nasal cavity)
- 7. Nasal concha
- 8. Pharynx
- 9. (Premaxillary bone)
- 10. Infraorbital sinus
- 11. Nasal septum
- 12. Choana
- 13. (Palatine bone)
- 14. (Sphenoid bone)
- 15. Jugal [zygomatic] bone
- 16. Frontal bone
- 17. Pterygoid bone
- 18. Eyeball

- 19. Scleral ossicle
- 20. (Interorbital septum)
 21. Lens of eyeball
 22. (Trachea)

- 23. (Cerebrum)
- 24. (External ear canal)
- 25. (Cerebellum)
- 26. (Spinal cord) 27. (Dens)
- 28. (Cere)
- 29. (Nare[s]) 30. (Feather)
- 31. (Pons)
- 32. (Occipital bone)
- 33. (Cervical vertebra)




Figure 16-17, I-L Type of Bird: Mallard Duck Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.4 kg Age: Adult

- 1. (Tongue)
- 2. Mandible
- 3. Hyoid bone
- 4. (Endotracheal tube)
- 5. (Keratinized maxillary beak)
- 6. (Nasal cavity)
- 7. (Nasal concha)
- 8. (Pharynx)
 9. (Premaxillary bone)
- 10. (Infraorbital sinus)
- 11. (Nasal septum)
- 12. (Choana)
- 13. (Palatine bone)
 14. (Sphenoid bone)
- 15. (Jugal [zygomatic] bone)
- 16. (Frontal bone)
- 17. (Pterygoid bone)
 18. (Eyeball)

- 19. (Scleral ossicle)
- 20. (Interorbital septum)
- 21. (Lens of eyeball)
 22. Trachea
- 23. Cerebrum
- 24. External ear canal
- 25. Cerebellum
- 26. Spinal cord 27. Dens
- 28. (Cere)
- 29. (Nare[s]) 30. (Feather)
- 31. (Pons)
- 32. (Occipital bone)
- 33. (Cervical vertebra)




Figure 16-17, M

Type of Bird: Mallard Duck Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.4 kg Age: Adult

- 1. (Tongue)
 2. (Mandible)
 3. (Hyoid bone) 4. (Endotracheal tube) 5. (Keratinized maxillary beak)
- 6. (Nasal cavity) (Nasal concha) 8. (Pharynx)
- 9. (Premaxillary bone) 10. (Infraorbital sinus)
- 11. (Nasal septum)
- 12. (Choana)
- 13. (Palatine bone)
- 14. (Sphenoid bone) 15. (Jugal [zygomatic] bone)
- 16. (Frontal bone)
- 17. (Pterygoid bone)
- 18. (Eyeball)

- 19. (Scleral ossicle)
- 20. (Interorbital septum)
- 21. (Lens of eyeball)
- 22. (Trachea)
- 23. (Cerebrum)
- 24. (External ear canal) 25. (Cerebellum)
- 26. Spinal cord 27. (Dens)
- 28. Cere
- 29. (Nare[s])
- 30. (Feather)
- 31. (Pons)
- 32. (Occipital bone)
- 33. (Cervical vertebra)




Figure 16-18, A-D

Type of Bird: Mallard Duck Type of Study: CT coelom Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.4 kg Age: Adult

- 1. Sternum
- 2. Thoracic air sac
- 3. Lung4. Esophagus
- 5. Thoracic vertebra
- 6. Heart
- 7. Blood vessels
- 8. (Thoracic extremity [wing])
- 9. Proventriculus
- 10. Ventriculus
- 11. Intestines
- 12. Pelvic extremity [leg]
 13. Abdominal air sac
 14. (Ilium)

- 15. (Liver)

- 16. (Kidney) 17. (Caudal vertebra)
- 18. (Rib)
- 19. (Aorta)
- 20. (Pubic bone)
- 21. (Cloaca)
- 22. (Clavicular air sac)
- 23. (Scapula)
- 24. (Syrinx)
- 25. (Ischium)
- 26. (Colon) 27. (Spinal cord)





Figure 16-18, E-HType of Bird: Mallard Duck Type of Study: CT coelom Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.4 kg Age: Adult

- 1. (Sternum)
- 2. (Thoracic air sac)
- 3. (Lung)
- 4. (Esophagus)
- 5. (Thoracic vertebra)
- 6. (Heart)
- (Blood vessels)
- 8. (Thoracic extremity [wing])
 9. (Proventriculus)
- 10. Ventriculus
- 11. Intestines
- 12. Pelvic extremity [leg]
- 13. Abdominal air sac 14. Ilium
- 15. Liver

- 16. Kidney17. Caudal vertebra
- 18. (Rib)
- 19. (Aorta)
- 20. (Pubic bone)
- 21. (Cloaca)
- 22. (Clavicular air sac)
- 23. (Scapula)
- 24. (Syrinx)
- 25. (Ischium)
- 26. (Colon) 27. (Spinal cord)

Red Tailed Hawk (Buteo jamaicensis)






Figure 17-1, A-B

Type of Bird: Red Tailed Hawk Type of Study: Viscera of the coelom
Projection: Laterolateral (right lateral
recumbency)
Weight of Bird: 1 kg

Gender: Unknown

Reproductive Status: Intact Age: Young adult

- 1. Trachea
- 2. (Crop)
- 3. Brachiocephalic artery and aorta
- 4. (Brachiocephalic artery)
- 5. Aorta
- 6. Pulmonary artery
- 7. Pulmonary vein
- 8. Heart
- 9. Left atrium
- 10. Esophagus
- 11. Liver
- 12. Lung
- 13. (Syrinx) 14. Gonad
- 15. (Ovary) 16. (Testes)

- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. Cloaca
- 21. (Cervical air sac)
 22. (Clavicular air sac)
- 23. Thoracic air sac
- 24. Abdominal air sac
- 25. (Apex of heart)
- 26. Interface between caudal thoracic and abdominal air sacs
- 27. Kidneys
- 28. Spleen




Figure 17-2, A-B Type of Bird: Red Tailed Hawk Type of Study: Viscera of the coelom Projection: Ventrodorsal Weight of Bird: 1 kg Gender: Unknown Reproductive Status: Intact Age: Young adult

- Trachea
 (Cervical air sac)
 Crop
- 4. Clavicular air sac
- 5. Brachiocephalic artery and aorta6. Heart base vessel7. Heart

- 8. Lung9. Thoracic air sac
- 10. Liver

- 11. Kidneys12. Proventriculus
- 13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. Cloaca




Figure 17-3, A-B

Type of Bird: Red Tailed Hawk Type of Study: Head

Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 1.2 kg Gender: Unknown Reproductive Status: Intact Age: Young adult

- 1. Keratinized maxillary beak
- 2. Keratinized mandibular beak
- 3. Premaxillary bone
- 4. Mandible
- 5. (Hyoid bones) 6. Palatine bone
- 7. (Pterygoid bone)
- 8. External nares
- 9. (Jugal [zygomatic] bone)
 10. Craniofacial flexion zone
- 11. Orbit
- 12. Frontal bone
- 13. Cranium
- 14. Temporal bone
- 15. (Quadrate bone)
 16. Parietal bone




Figure 17-4, A-B
Type of Bird: Red Tailed Hawk
Type of Study: Head Projection: Ventrodorsal Weight of Bird: 1.2 kg Gender: Unknown Reproductive Status: Intact Age: Young adult

- 1. Keratinized maxillary beak
- 2. Premaxillary bone
 3. Mandible
 4. Orbit

- 5. Jugal [zygomatic] bone
- 6. Scleral ossicles7. (Quadrate bone)
- 8. Cranium




Figure 17-5, A-B

Type of Bird: Red Tailed Hawk Type of Study: Whole body skeleton Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 1 kg

Gender: Unknown

Reproductive Status: Intact

Age: Young adult

- 1. Cervical vertebra
- 2. Clavicle
- 3. Humerus
- 4. Coracoid
- 5. Scapula6. Rib
- 7. Uncinate process of rib 8. Thoracic vertebra
- 9. Synsacrum
- 10. Ilium
- 11. Head of femur
- 12. Ilioischiadiac foramen
- 13. Ischium

- 14. Pubis
- 15. Femur
- 16. (Obturator foramen)
- 17. Patella
- 18. Tibiotarsal bone
- 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae 22. Pygostyle




Figure 17-6, A-B Type of Bird: Red Tailed Hawk Type of Study: Whole body skeleton Projection: Ventrodorsal Weight of Bird: 1 kg Gender: Unknown Reproductive Status: Intact Age: Young adult

- 1. Cervical vertebra
- Clavicle
 Humerus
- 4. Coracoid
- 5. Scapula
- 6. Rib7. Uncinate process of rib
- 8. Thoracic vertebra
- 9. Synsacrum
- 10. Ilium
- 11. Head of femur
- 12. (Ilioischiadiac foramen)
- 13. İschium
- 14. Pubis 15. Femur

- 16. (Obturator foramen)
- 17. (Patella)
- 18. Tibiotarsal bone
- 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae
- 22. Pygostyle
- 23. Apex carinae
- 24. Dorsal tubercle of humerus25. Head of humerus
- 26. Ventral tubercle of humerus
- 27. Trochanter of femur




Figure 17-7, A-B
Type of Bird: Red Tailed Hawk
Type of Study: Wing Projection: Mediolateral Weight of Bird: 1 kg Gender: Unknown Reproductive Status: Intact Age: Young adult

- Head of humerus
 Dorsal tubercle of humerus
 Dorsal condyle of humerus
- 4. Radius
- 5. Radial carpal bone
- 6. Alula
 7. Major metacarpal bone
 8. Phalanges of major digit

- 9. Ventral tubercle of humerus
- 10. Humerus
- 11. Ventral condyle of humerus
- 12. Ulna

- 13. Ulnar carpal bone14. Minor metacarpal bone15. Phalanges of minor digit




Figure 17-8, A-B
Type of Bird: Red Tailed Hawk
Type of Study: Wing
Projection: Caudocranial
Weight of Bird: 1 kg Gender: Unknown Reproductive Status: Intact Age: Young adult

- 1. Head of humerus
- 2. Ventral tubercle of humerus
- 3. Ventral condyle of humerus
- 4. Radius
 5. Radial carpal bone
 6. Alula
- 7. Minor metacarpal bone
- 8. Phalanges of major digit
- 9. Dorsal tubercle of humerus
- 10. Humerus
- 11. Dorsal condyle of humerus
- 12. Ulna
- 13. Ulnar carpal bone
- 14. Major metacarpal bone
- 15. Phalanges of minor digit




Figure 17-9, A-B Type of Bird: Red Tailed Hawk Type of Study: Pelvic limb Projection: Craniocaudal Weight of Bird: 1.2 kg Gender: Unknown Reproductive Status: Intact Age: Young adult

- 1. Trochanter of femur
- 2. Femur
- 3. Lateral condyle of femur
- 4. Fibula
- 5. Lateral condyle of tibiotarsal bone
- 6. Tarsometatarsal bone
- 7. Trochlea of tarsometatarsal bone 8. Digit IV
- 9. Digit III
- 10. Head of femur

- 11. Medial condyle of femur12. Tibiotarsal bone
- 13. Medial condyle of tibiotarsal bone
- 14. Digit I
- 15. Digit II 16. Phalanges
- 17. Patella
- 18. Hypotarsal crest of tarsometatarsal bone
- 19. Metatarsal I




Figure 17-10, A-B

Type of Bird: Red Tailed Hawk Type of Study: Pelvic limb Projection: Mediolateral Weight of Bird: 1.2 kg Gender: Unknown Reproductive Status: Intact Age: Young adult

- 1. Femur
- 2. Patella
 3. Condyles of femur
 4. Fibula
 5. Tibiotarsal bone

- 6. Condyles of tibiotarsal bone7. Hypotarsal crest of tarsometatarsal bone
- 8. Tarsometatarsal bone
- 9. Trochlea of tarsometatarsal bone

- 10. Digit I 11. Digit II 12. Digit III
- 13. Digit IV
- 14. Phalanges 15. Metatarsal I




Figure 17-11, A-B

Type of Bird: Red Tailed Hawk
Type of Study: Distal pelvic limb
Projection: Mediolateral
Weight of Bird: 1.2 kg
Gender: Unknown
Reproductive Status: Intact
Age: Young adult

- 1. Tarsometatarsal bone
- 2. Trochlea of tarsometatarsal bone
- 3. Metatarsal bone I
- 4. Digit I
- 5. Digit II
- 6. Digit III 7. Digit IV

- 8. Phalanges
- 9. (Hypotarsal crest of tarsometatarsal bone)
- 10. (Condyles of tibiotarsal bone)

NOTE: Structures in parentheses are not labeled.




Figure 17-12, A-B

Type of Bird: Red Tailed Hawk Type of Study: Distal pelvic limb Projection: Dorsoplantar Weight of Bird: 1.2 kg Gender: Unknown Reproductive Status: Intact Age: Young adult

- 1. Tarsometatarsal bone
- 2. Trochlea of tarsometatarsal bone
- 3. Metatarsal bone I
- 4. Digit I
- 5. Digit II
- 6. Digit III 7. Digit IV

- 8. Phalanges
- 9. (Hypotarsal crest of tarsometatarsal bone)
- 10. (Condyles of tibiotarsal bone)




Figure 17-13, A
Type of Bird: Red Tailed Hawk Type of Study: Gastrointestinal positive contrast study

Contrast Study
Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 20 ml administered via gavage tube Projection: Laterolateral (right lateral

recumbency)
Weight of Bird: 1.2 kg
Gender: Unknown
Reproductive Status: Intact Age: Young adult

lmage	Time (hr)
A	Scout

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)
- 8. (Cloaca)
- 9. (Proventricular-ventricular isthmus)




Figure 17-13, B-C
Type of Bird: Red Tailed Hawk
Type of Study: Gastrointestinal positive contrast study

Contrast study
Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)

20 ml administered via gavage tube Projection: Laterolateral (right lateral recumbency) Weight of Animal: 1.2 kg

Gender: Unknown Reproductive Status: Intact

Age: Young adult

lmage	Time (hr)
В	0.25
С	1.0

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus
 5. Intestines

- 6. (Duodenum)7. Large intestine
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 17-13, D-E
Type of Bird: Red Tailed Hawk
Type of Study: Gastrointestinal positive contrast study

Contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)

20 ml administered via gavage tube
Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 1.2 kg
Gender: Unknown Reproductive Status: Intact Age: Young adult

lmage	Time (hr)
D	3.0
E.	5.0

- Crop
 Esophagus
- 3. Proventriculus
- 4. Ventriculus

- 5. Intestines
 6. (Duodenum)
 7. Large intestine
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 17-14, A-B

Type of Bird: Red Tailed Hawk Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 20 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 1.2 kg Gender: Unknown Reproductive Status: Intact Age: Young adult

lmage	Time (hr)
A	Scout
В	0.25

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. (Large intestine)
 8. (Cloaca)
- 9. (Proventricular-ventricular isthmus)




Figure 17-14, C-DType of Bird: Red Tailed Hawk

Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutici, Inc., Lafayette, IN) 20 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 1.2 kg Gender: Unknown Reproductive Status: Intact Age: Young adult

lmage	Time (hr)
С	1.0
D	3.0

- Crop
 Esophagus
- 3. Proventriculus4. Ventriculus
- 5. Intestines 6. Duodenum
- 7. Large intestine
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 17-14, E

Type of Bird: Red Tailed Hawk Type of Study: Gastrointestinal positive contrast study
Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/r; Lafayette

Pharmaceutical, Inc., Lafayette, IN) 20 ml administered via gavage tube Projection: Ventrodorsal Weight of Bird: 1.2 kg

Gender: Unknown Reproductive Status: Intact Age: Young adult

lmage	Time (hr)
Е	5.0

- Crop
 (Esophagus)
 Proventriculus
 Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. Large intestine
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 17-15, A-DType of Bird: Red Tailed Hawk Type of Study: CT coelom Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.1 kg Age: Adult

- 2. (Thoracic air sac)
- 3. Lung
- 4. (Esophagus)
- 5. (Thoracic vertebra)
- 6. Heart
- 7. Blood vessels
- 8. Thoracic extremity (wing)
- 9. (Proventriculus)
 10. (Ventriculus)
- 11. (Intestines)
- 12. (Pelvic extremity (leg))
- 13. (Abdominal air sac) 14. (Ilium) 15. (Liver)

- 18. (Rib)
 19. (Aorta)
 20. (Pubic bone)
- 21. (Cloaca)
- 22. (Clavicular air sac)
- 23. Scapula
- 24. (Syrinx)
- 25. (Ischium)
- 26. (Colon)
- 27. (Spinal cord)




Figure 17-15, E-H

Type of Bird: Red Tailed Hawk Type of Study: CT coelom Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.1 kg Age: Adult

- $1. \ Sternum$
- 2. (Thoracic air sac)
- 3. Lung
- 4. (Esophagus)
- 5. (Thoracic vertebra)
- 6. Heart
- 7. Blood vessels
- 8. (Thoracic extremity (wing))
- 9. Proventriculus
 10. (Ventriculus)
- 11. (Intestines)
- 12. (Pelvic extremity (leg))
- 13. (Abdominal air sac)
 14. (Ilium)
- 15. Liver

- 16. (Kidney) 17. (Caudal vertebra)

- 18. (Rib)
 19. (Aorta)
 20. (Pubic bone)
- 21. (Cloaca)
- 22. (Clavicular air sac)
- 23. (Scapula) 24. (Syrinx)
- 25. (Ischium) 26. (Colon)
- 27. (Spinal cord)




Figure 17-15, I-L

Type of Bird: Red Tailed Hawk Type of Study: CT coelom Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.1 kg Age: Adult

- 1. (Sternum)
- 2. (Thoracic air sac)
- 3. (Lung)
- 4. (Esophagus) 5. Thoracic vertebra
- 6. (Heart)
- 7. (Blood vessels)
- 8. (Thoracic extremity (wing))
 9. Proventriculus
- 10. Ventriculus
- 11. Intestines
- 12. Pelvic extremity (leg)
- 13. (Abdominal air sac)
- 14. (Ilium)
- 15. (Liver)

- 16. Kidney
- 17. (Caudal vertebra)
- 18. (Rib)
- 19. (Aorta)
- 20. (Pubic bone)
- 21. (Cloaca)
- 22. (Clavicular air sac)
- 23. (Scapula)
- 24. (Syrinx)
- 25. (Ischium)
- 26. (Colon)
- 27. (Spinal cord)




Figure 17-15, M

Type of Bird: Red Tailed Hawk Type of Study: CT coelom Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.1 kg Age: Adult

1. (Sternum)

2. (Thoracic air sac)

3. (Lung)

4. (Esophagus)

5. (Thoracic vertebra) 6. (Heart)

7. (Blood vessels)

8. (Thoracic extremity (wing))
9. (Proventriculus)

10. (Ventriculus)

11. (Intestines)

12. Pelvic extremity (leg)

13. (Abdominal air sac)
14. (Ilium)

15. (Liver)

16. (Kidney) 17. Caudal vertebra

18. (Rib)

19. (Aorta) 20. (Pubic bone)

21. Cloaca

22. (Clavicular air sac)
23. (Scapula)


24. (Syrinx)

25. (Ischium)

26. (Colon)

27. (Spinal cord)

Great Horned Owl (Bubo virginianus)






Figure 18-1, A-B

Type of Bird: Great Horned Owl Type of Study: Viscera of the coelom Projection: Laterolateral (right lateral

recumbency)
Weight of Bird: 1.3 kg Gender: Unknown Reproductive Status: Intact

Age: Adult

- 1. Trachea
- 2. (Crop)
- 3. (Brachiocephalic artery and aorta)
- 4. Brachiocephalic artery
- 5. Aorta
- 6. Pulmonary artery
- 7. Pulmonary vein
- 8. Heart
- 9. Left atrium
- 10. Esophagus
- ll. Liver
- 12. Lung
- 13. (Syrinx) 14. (Gonad) 15. (Ovary) 16. (Testes)

- 17. Proventriculus
- 18. Ventriculus
- 19. Intestines
- 20. (Cloaca)
 21. Cervical air sac
- 22. (Clavicular air sac)
- 23. (Thoracic air sac)
- 24. (Abdominal air sac) 25. (Apex of heart)
- 26. Interface between caudal thoracic and abdominal air sacs
- 27. Kidneys
- 28. Spleen




Figure 18-2, A-BType of Bird: Great Horned Owl Type of Study: Viscera of the coelom Projection: Ventrodorsal Weight of Bird: 1.3 kg Gender: Unknown Reproductive Status: Intact Age: Adult

- Trachea
 (Cervical air sac)
 (Crop)
 Claudiular air sac

- 5. Brachiocephalic artery and aorta6. Heart base vessel7. Heart

- 8. Lung
 9. Thoracic air sac
 10. Liver

- 11. Kidneys12. (Proventriculus)
- 13. Ventriculus
- 14. Intestines
- 15. Abdominal air sac
- 16. Cloaca




Figure 18-3, A-B

Type of Bird: Great Horned Owl Type of Study: Head

Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 1.3 kg

Weight of Bird: 1.3 kg Gender: Unknown Reproductive Status: Intact

Age: Adult

- 1. Keratinized maxillary beak
- 2. Keratinized mandibular beak
- 3. Premaxillary bone
- 4. Mandible
- 5. (Hyoid bones)
- 6. (Palatine bone)
- 7. (Pterygoid bone)
- 8. External nares
- 9. (Jugal [zygomatic] bone)
- 10. Craniofacial flexion zone
- 11. Orbit
- 12. Frontal bone
- 13. Cranium
- 14. Temporal bone
- 15. (Quadrate bone)
- 16. Parietal bone




Figure 18-4, A-B
Type of Bird: Great Horned Owl
Type of Study: Head
Projection: Ventrodorsal
Weight of Bird: 1.3 kg
Gender: Unknown
Reproductive Status: Intact
Age: Adult

- 1. Keratinized maxillary beak
- Premaxillary bone
 Mandible
- 4. Orbit
- 5. Jugal [zygomatic] bone
- 6. (Scleral ossicles)
- 7. Quadrate bone 8. Cranium




Figure 18-5, A-B

Type of Bird: Great Horned Owl
Type of Study: Whole body skeleton
Projection: Laterolateral (right lateral recumbency)
Weight of Bird: 1.3 kg

Gender: Unknown Reproductive Status: Intact

Age: Adult

- 4. Coracoid
- 5. Scapula
- 6. Rib
- 7. Uncinate process of rib
- 8. Thoracic vertebra
- 9. Synsacrum
- 10. Ilium
- 11. Head of femur

- 13. Ischium
- 14. Pubis
- 15. Femur
- 16. Obturator foramen
- 17. Patella
- 18. Tibiotarsal bone
- 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae
- 22. Pygostyle




Figure 18-6, A-B
Type of Bird: Great Horned Owl Type of Study: Whole body skeleton Projection: Ventrodorsal Weight of Bird: 1.3 kg Gender: Unknown Reproductive Status: Intact Age: Adult

- 1. Cervical vertebra
- 2. Clavicle
- 3. (Humerus)

- 4. Coracoid
 5. Scapula
 6. Rib
 7. (Uncinate (Uncinate process of rib)
- 8. Thoracic vertebra
- 9. Synsacrum
- 10. Ilium
- 11. Head of femur
- 12. Ilioischiadiac foramen
- 13. Ischium
- 14. Pubis 15. Femur

- 16. (Obturator foramen)17. (Patella)
- 18. Tibiotarsal bone
- 19. Fibula
- 20. Sternum
- 21. Caudal vertebrae
- 22. Pygostyle
- 23. Apex carinae
- 24. Dorsal tubercle of humerus25. Head of humerus
- 26. Ventral tubercle of humerus
- 27. Trochanter of femur




Figure 18-7, A-B Type of Bird: Great Horned Owl Type of Study: Wing Projection: Mediolateral Weight of Bird: 1.3 kg

Gender: Unknown Reproductive Status: Intact Age: Adult

- 1. Head of humerus
- Dorsal tubercle of humerus
 Dorsal condyle of humerus
- 4. Radius
- 5. Radial carpal bone
- 6. Alula
- 7. Major metacarpal bone8. Phalanges of major digit

- 9. Ventral tubercle of humerus
- 10. Humerus11. Ventral condyle of humerus
- 12. Ulna
- 13. Ulnar carpal bone
- 14. Minor metacarpal bone 15. Phalanges of minor digit





Figure 18-8, A-B
Type of Brid: Great Horned Owl
Type of Study: Wing
Projection: Caudocranial
Weight of Bird: 1.3 kg Gender: Unknown Reproductive Status: Intact Age: Adult

- 1. Head of humerus
- 2. Ventral tubercle of humerus3. Ventral condyle of humerus

- 4. Radius5. Radial carpal bone
- 6. Alula
- 7. Minor metacarpal bone
- 8. Phalanges of major digit 9. (Dorsal tubercle of humerus)
- 10. Humerus
- 11. Dorsal condyle of humerus
- 12. Ulna
- 13. Ulnar carpal bone
- 14. Major metacarpal bone
- 15. Phalanges of minor digit



Type of Bird: Great Horned Owl Type of Study: Pelvic limb

Projection: Mediolateral Weight of Bird: 1.3 kg Gender: Unknown Reproductive Status: Intact

Age: Adult

- 3. Condyles of femur

- 5. Condyles of Telliul4. Fibula5. Tibiotarsal bone6. Condyles of tibiotarsal bone7. Hypotarsal crest of tarsometatarsal bone
- 9. Trochlea of tarsometatarsal bone

- 10. Digit I 11. Digit II 12. Digit III
- 13. Digit IV
- 14. Phalanges
- 15. Metatarsal I




Figure 18-10, A-B

Type of Bird: Great Horned Owl Type of Study: Pelvic limb Projection: Craniocaudal Weight of Bird: 1.3 kg Gender: Unknown Reproductive Status: Intact Age: Adult

- 1. Trochanter of femur
- 2. Femur3. Lateral condyle of femur
- 4. Fibula
- 5. Lateral condyle of tibiotarsal bone6. Tarsometatarsal bone
- 7. Trochlea of tarsometatarsal bone
- 8. Digit IV
- 9. Digit III 10. Head of femur
- 11. Medial condyle of femur

- 12. Tibiotarsal bone
- 13. Medial condyle of tibiotarsal bone
- 14. Digit I
- 15. Digit II 16. Phalanges 17. (Patella)
- 18. (Hypotarsal crest of tarsometatarsal bone)
- 19. Metatarsal I




Figure 18-11, A-B
Type of Bird: Great Horned Owl
Type of Study: Distal pelvic limb
Projection: Mediolateral
Weight of Bird: 1.3 kg
Gender: Unknown
Pennsylvative Status: Intest

Reproductive Status: Intact

Age: Adult

- 1. Tarsometatarsal bone
- 2. Trochlea of tarsometatarsal bone
- 3. Metatarsal bone I

- 4. Digit I
 5. Digit II
 6. Digit III

- 7. Digit IV8. Phalanges
- 9. Hypotarsal crest of tarsometatarsal bone
 10. Condyles of tibiotarsal bone




Figure 18-12, A-B
Type of Bird: Great Horned Owl
Type of Study: Distal pelvic limb Projection: Dorsoplantar Weight of Bird: 1.3 kg Gender: Unknown Reproductive Status: Intact Age: Adult

- Tarsometatarsal bone
 Trochlea of tarsometatarsal bone
 Metatarsal bone I

- 4. Digit I
 5. Digit II
 6. Digit III
 7. Digit IV

- 8. Phalanges9. (Hypotarsal crest of tarsometatarsal bone)
- 10. (Condyles of tibiotarsal bone)




Figure 18-13, A-B

Type of Bird: Great Horned Owl Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN)
23 ml administered via gavage tube
Projection: Laterolateral (right lateral

recumbency) Weight of Bird: 1.3 kg Gender: Unknown Reproductive Status: Intact Age: Adult

Time (hr) Image A Scout 0.25 В

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 18-13, C-D
Type of Bird: Great Horned Owl
Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 23 ml administered via gavage tube Projection: Laterolateral (right lateral recumbency)

Weight of Bird: 1.3 kg Gender: Unknown Reproductive Status: Intact Age: Adult

Image	Time (hr)
С	1.0
D	3.0

- Crop
 Esophagus
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. (Large intestine)
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 18-13, E

Type of Bird: Great Horned Owl Type of Study: Gastrointestinal positive contrast study

Contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 23 ml administered via gavage tube Projection: Laterolateral (right lateral

recumbency)

Weight of Bird: 1.3 kg Gender: Unknown Reproductive Status: Intact

Age: Adult

lmage	Time (hr)
Е	5.0

- 1. Crop
- 2. Esophagus3. Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. Duodenum
- 7. Large intestine
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 18-14, A-B

Type of Bird: Great Horned Owl Type of Study: Gastrointestinal positive contrast study

Contrast Study
Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 23 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 1.3 kg Gender: Unknown Reproductive Status: Intact

Age: Adult

Image	Time (hr)
A	Scout
В	0.25

- 1. Crop
- 2. (Esophagus)
- 3. Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
 7. Large intestine
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 18-14, C-D

Type of Bird: Great Horned Owl Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 23 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 1.3 kg Gender: Unknown Reproductive Status: Intact Age: Adult

lmage	Time (hr)
С	1.0
D	3.0

- Crop
 (Esophagus)
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. Large intestine
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 18-14, E

Type of Bird: Great Horned Owl Type of Study: Gastrointestinal positive contrast study

Contrast Medium: Barium sulfate suspension (Novopaque® 30% w/v; Lafayette Pharmaceutical, Inc., Lafayette, IN) 23 ml administered via gavage tube

Projection: Ventrodorsal Weight of Bird: 1.3 kg Gender: Unknown Reproductive Status: Intact Age: Adult

lmage	Time (hr)
Е	5.0

- Crop
 (Esophagus)
 Proventriculus
- 4. Ventriculus
- 5. Intestines
- 6. (Duodenum)
- 7. Large intestine
- 8. Cloaca
- 9. (Proventricular-ventricular isthmus)




Figure 18-15, A-D Type of Bird: Great Horned Owl Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.4 kg Age: Adult

- 3. (Hyoid bone) 4. Endotracheal tube 5. Keratinized maxillary beak 6. (Nasal cavity) 7. Nasal concha 8. (Pharynx) 9. (Premaxillary bone)
- 12. (Choana)
 13. (Palatine bone) 14. (Sphenoid bone)
- 16. (Frontal bone)
 17. (Pterygoid bone)
 18. Eyeball

- 1. (Tongue) 2. Mandible

- 10. (Infraorbital sinus)
- 11. Nasal septum
- 15. (Jugal [zygomatic] bone)

- 19. (Scleral ossicle)
- 20. (Interorbital septum)
 21. (Lens of eyeball)
 22. (Trachea)

- 23. (Cerebrum)
- 24. (External ear canal)
- 25. (Cerebellum)
- 26. (Spinal cord) 27. (Dens)
- 28. (Cere)
- 29. (Nare[s]) 30. (Feather)
- 31. (Pons)
- 32. (Occipital bone)
- 33. Cervical vertebra




Figure 18-15, E-H Type of Bird: Great Horned Owl Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.4 kg Age: Adult

- 1. Tongue
- 2. Mandible3. Hyoid bone
- 4. (Endotracheal tube)
- 5. (Keratinized maxillary beak)
- 6. Nasal cavity
- 7. Nasal concha 8. (Pharynx)
- 9. (Premaxillary bone)
- 10. Infraorbital sinus
- 11. Nasal septum
- 12. Choana
- 13. (Palatine bone)
- 14. (Sphenoid bone)
- 15. Jugal [zygomatic] bone 16. (Frontal bone) 17. (Pterygoid bone)

- 18. Eyeball

- 19. (Scleral ossicle)20. Interorbital septum21. (Lens of eyeball)
- 22. (Trachea)
- 23. Cerebrum
- 24. (External ear canal)
 25. (Cerebellum)
- 26. (Spinal cord) 27. (Dens)

- 28. (Cere) 29. (Nare[s])
- 30. (Feather)
- 31. (Pons)
- 32. (Occipital bone)
- 33. Cervical vertebra




Figure 18-15, I-L Type of Bird: Great Horned Owl Type of Study: CT head Contrast Medium: None Imaging Plane: Transverse Weight of Bird: 1.4 kg Age: Adult

- 4. (Endotracheal tube) 5. (Keratinized maxillary beak) 6. (Nasal cavity)
 7. (Nasal concha) 8. (Pharynx) 9. (Premaxillary bone) 10. (Infraorbital sinus) 11. (Nasal septum)
 12. (Choana)
 13. (Palatine bone)
- 15. (Jugal [zygomatic] bone) 16. (Frontal bone)
 17. (Pterygoid bone)
 18. (Eyeball)

- (Tongue)
 Mandible
- 3. Hyoid bone

- 14. (Sphenoid bone)

- 19. (Scleral ossicle)
- 20. (Interorbital septum)
 21. (Lens of eyeball)
- 22. (Trachea)
- 23. Cerebrum
- 24. (External ear canal)
 25. Cerebellum
 26. (Spinal cord)
 27. (Dens)
- 28. (Cere)
- 29. (Nare[s]) 30. Feather
- 31. Pons
- 32. (Occipital bone)
- 33. Cervical vertebra

INDEX

A	Blue and gold macaw (Ara ararauna) (Continued)
African grey parrot (Psittacus erithacus erithacus), 67-73	laterolateral view of, 130f
infraorbital sinus contrast study	ventrodorsal view of, 131f
laterolateral view of, 72f	viscera of coelom
ventrodorsal view of, 73f	laterolateral view of, 126f
viscera of coelom	ventrodorsal view of, 127f
laterolateral view of, 68f	wing
ventrodorsal view of, 69f	caudocranial view, 133f
whole body skeleton	mediolateral view, 132 <i>f</i>
laterolateral view of, 70f	Bones. See Head; Pelvic limb; Skeleton, whole body; Wing
ventrodorsal view of, 71f	Bubo virginianus. See Great horned owl.
Agapornis roseicollis. See Peach-faced lovebird.	Budgerigar (Melopsittacus undulates), 25-27
Air sacs	viscera of coelom
See also Viscera.	laterolateral view of, 26f
inflation of, 3	ventrodorsal view of, 27f
Airway aspiration during radiography, 3-4	Buteo jamaicensis. See Red tailed hawk.
Alula. See Wing.	Butto junimeerisis. See Red tailed Hawk.
Amazona amazonica. See Orange-winged Amazon parrot.	C
Anas platyrhynchos. See Mallard duck.	Cacatua (Cacatua) alba. See Umbrella cockatoo.
Anatomic art, 17-23	Cacatua galerita. See Sulphur-crested cockatoo.
lateral view	Cacatua (Licmetis) goffini). See Goffin cockatoo.
of skeleton, 22f	Cacatua moluccensis. See Moluccan cockatoo.
of skull, 20f	Carbon dioxide for contrast medium removal, 13
of viscera, 18f	Caudocranial view
ventral view	of pelvic limb
of skeleton, 23f	of great horned owl, 259f
of skull, 21 <i>f</i>	of mallard duck, 213f
of viscera, 19f	of Moluccan cockatoo, 171f
Anatomy of the Domestic Birds, 2	of orange-winged Amazon parrot, 89f
Anesthesia	of pigeon, 189f
for digestive tract contrast studies, 10	of red tailed hawk, 236f
for radiography, 3-4, 6	positioning techniques
Ara ararauna. See Blue and gold macaw.	of leg, 7, 12f
Aratinga solstitialis. See Sun conure.	of wing, 6-7, 10f
Arteries. See Viscera.	of wing
Asymetrix Detail Intensifying Screen, 2	of blue and gold macaw, 133f
Atlas of Avian Anatomy: Osteology-Arthrology-Mycology, 2	of great horned owl, 257f
Atlas of Avian Radiographic Anatomy, 2	of mallard duck, 211f
Atlas of Radiographic Anatomy and Diagnosis of Cage	of Moluccan cockatoo, 169f
Birds, 2	of orange-winged Amazon parrot, 87f
Birtis, Z	of pigeon, 187f
В	of red tailed hawk, 235f
Bandage gauze, in positioning, 5, 7	Cerebellum. See Head.
Barium sulfate as contrast medium, 9	Cerebrum. See Head.
Beak, maxillary and mandibular. See Head.	Clavicle. See Skeleton, whole body.
Bird Board, 4, 4f	Cloaca
Blue and gold macaw (Ara ararauna), 125-145	See also Viscera.
CT head, transverse view of, 142-145f	double contrast studies of, 13
gastrointestinal positive contrast study	laterolateral view
laterolateral view of, 136-138f	of orange-winged Amazon parrot, 102f-103f
ventrodorsal view of, 139-141f	of pigeon, 200f
head	ventrodorsal view
laterolateral view of, 128f	of orange-winged Amazon parrot, 104 <i>f</i> -105 <i>f</i>
ventrodorsal view of, 129f	of pigeon, 201f
pelvic limb, mediolateral view of, 134f	Cockatiel (Nymphicus hollandicus), 39-45
skeleton, whole body	gastrointestinal positive contrast study
merculi, militie body	5 de la contra postar e contras suas

Cockatiel (Nymphicus hollandicus) (Continued)	Foot, mediolateral and dorsoplantar studies of, 7-8, 13, 14f
ventrodorsal view of, 42f-43f, 44f-45f	Foramina. See Skeleton, whole body.
viscera of coelom	
laterolateral view of, 40f	G
ventrodorsal view of, 41f	Gastrografin, 10
Cockatoo. See Goffin cockatoo; Moluccan cockatoo;	Gastrointestinal tract
Sulphur-crested cockatoo; Umbrella cockatoo.	distension during radiography, 3-4
Coelom	double contrast study
See also Viscera of coelom.	laterolateral view
CT transverse view	of orange-winged Amazon parrot, 98f-99f
of mallard duck, 225f-226f	of Senegal parrot, 62f-63f ventrodorsal view of orange-winged Amazon parrot,
of red tailed hawk, 245f-248f	100f- $101f$
Color Atlas of Avian Anatomy, 2 Columba livia. See Pigeon.	positive contrast study
Computerized tomography (CT), 14-15	laterolateral view
transverse view	of blue and gold macaw, 136f-138f
coelom	of great horned owl, 262f-264f
of mallard duck, 225 <i>f</i> -226 <i>f</i>	of mallard duck, 216f
of orange-winged Amazon parrot, 113f-117f	of Moluccan cockatoo, 174f
of red tailed hawk, 245f-248f	of orange-winged Amazon parrot, 92f-94f
head	of peach-faced lovebird, 32f-34f
of great horned owl, 268f-270f	of pigeon, 184 <i>f</i> -186 <i>f</i>
of mallard duck, 221 <i>f</i> -224 <i>f</i>	of red tailed hawk, 239f-241f
of orange-winged Amazon parrot, 110f-112f	of Senegal parrot, 54f-57f
of sulphur-crested cockatoo, 152f, 153f	ventrodorsal view
Contrast studies	of blue and gold macaw, 139f-141f
of gastrointestinal tract, 8-14	of cockatiel, 42f-45f
of urinary tract, 14	of great horned owl, 265 <i>f</i> -267 <i>f</i> of mallard duck, 217 <i>f</i> -218 <i>f</i>
Coracoid. See Skeleton, whole body. Craniofacial flexion zone. See Head.	of Moluccan cockatoo, 175
Cranium. See Head.	of orange-winged Amazon parrot, 95f-96f
Crop. See Viscera.	of peach-faced lovebird, 35f-37f
CT. See Computerized tomography.	of pigeon, 197-199
on our compactified tomography.	of red tailed hawk, 242-244
D	of Senegal parrot, 58f-61f
Diatrizoate meglumine, 10	radiographic contrast studies of, 8-14
Digestive tract. See Gastrointestinal tract.	Gavage tube, 10
Digital radiology systems, 3	Goffin cockatoo (Cacatua [Licmetis] goffini), 147-149
Digits of foot, positioning of, 8	viscera of coelom
Dorsoplantar view of pelvic limb	laterolateral view of, 148f
of great horned owl, 261f	ventrodorsal view of, 149f
of mallard duck, 215f	Gonads. See Viscera.
of Moluccan cockatoo, 173f	Great horned owl (Bubo virginianus), 249-270
of orange-winged Amazon parrot, 91f	gastrointestinal positive contrast study laterolateral view of 262 <i>f</i> -264 <i>f</i>
of pigeon, 191 <i>f</i> positioning techniques for, 7-8, 13 <i>f</i> , 14 <i>f</i>	ventrodorsal view of, 265f-267f
of red tailed hawk, 238f	head, CT transverse view, 268f-270f
Dose of contrast medium, 10	laterolateral view of, 252f
Double contrast digestive tract studies, 12-13	ventrodorsal view of, 253f
Duck. See Mallard duck.	pelvic limb
7	caudocranial view of, 259f
E	dorsoplantar view of, 261f
Equipment for radiology. See Radiology, equipment for.	mediolateral view of, 258f, 260f
Esophagus. See Viscera.	skeleton, whole body
Excretory urogram	laterolateral view of, 254f
laterolateral view	ventrodorsal view of, 255 <i>f</i>
of mallard duck, 219 <i>f</i> , 220 <i>f</i>	viscera of coelom
of orange-winged Amazon parrot, 106f-107f ventrodorsal view of orange-winged Amazon parrot, 108f-109f	laterolateral view of, 250
Exposure factors, timing of, 2, 3, $3t$	ventrodorsal view of, 251f
Eyeball. See Head.	wing caudocranial view of, 257f
Dyeban. See Flead.	mediolateral view of, 256f
F	Guillotine, in positioning, 5-7
Fasting period, 8	duniotine, in positioning, 5 7
Feather. See Head.	Н
Feeding before radiography, 3, 9	Hawk, red tailed. See Red tailed hawk.
Femur. See Skeleton, whole body.	Head
Fibula. See Skeleton, whole body.	CT transverse view
Film, radiographic, 2-3	of blue and gold macaw, 142f-145f
Flight feathers, positioning of, 5	of great horned owl, 268f-270f
Focal film distance (FFD), 2	of mallard duck, 221f-224f
Focal spot, 2	of Moluccan cockatoo, 176f-178f
Food deprivation, 3	of sulphur-crested cockatoo, 152f, 153f

Head (Continued)	Laterolateral view (Continued)
laterolateral and ventrodorsal studies, 4, 5f, 6f	of coelom, 4-5, 7 <i>f</i>
laterolateral view	of head, 4, 5 <i>f</i> , 6 <i>f</i>
of blue and gold macaw, 128f	respiratory series of pigeon, 192f-193f
of great horned owl, 252f	of skeleton, whole body
of mallard duck, 206f	of African grey parrot, 70f
of Moluccan cockatoo, 164f	of blue and gold macaw, 130f
of orange-winged Amazon parrot, 80f	of great horned owl, 254f
of pigeon, 182f	of mallard duck, 208f
of red tailed hawk, 230f	of Moluccan cockatoo, 166f
oblique view of orange-winged Amazon parrot, 81f	of orange-winged Amazon parrot, 84f
ventrodorsal view	of pigeon, 184f
of blue and gold macaw, 129f	of red tailed hawk, 232f
of great horned owl, 253f	of viscera of coelom
of mallard duck, 207f	of African grey parrot, 68f
of Moluccan cockatoo, 165f	of blue and gold macaw, 126f
of orange-winged Amazon parrot, 82f-83f	of budgerigar, 26f
of pigeon, 183f	of cockatiel, 40f
of red tailed hawk, 231f	of Goffin cockatoo, 148
Heart. See Viscera.	of great horned owl, 250f
Humerus. See Skeleton, whole body; Wing.	of mallard duck, 204f
Hyperalimentation preparations, 9	of Moluccan cockatoo, 160f, 162f
Ī	of orange-winged Amazon parrot, 76f, 78f
Ilium. See Skeleton, whole body.	of peach-faced lovebird, 30f
Infraorbital sinus contrast study	of pigeon, 180f
of African grey parrot	of red tailed hawk, 228f
laterolateral view of, 72 <i>f</i>	of Senegal parrot, 52f
ventrodorsal view of, 73f	of sun conure, 48 <i>f</i> of umbrella cockatoo, 156 <i>f</i>
MRI studies, 15	Leg
Inglovies (crop), emptying of, 8	See also Pelvic limb.
Innovet Select 20kHz High-Frequency Radiographic Machine, 2	craniocaudal study of, 7, 12f
Intestines. See Viscera.	mediolateral study of, 7, 11f
Ischium. See Skeleton, whole body.	Liver
	See also Viscera.
K	ultrasound of orange-winged Amazon parrot,
Kidney. <i>See</i> Viscera.	124f
Killivolt settings, 2	Lovebird, peach-faced. See Peach-faced lovebird.
	Lungs
L	See also Viscera.
Lateral view	aspiration of contrast medium, 12
of skeleton, 22f	CT studies of, 15
of skull, 20f	
of viscera, 18f	M
Laterolateral view	Macaw. See Blue and gold macaw.
of cloaca	Magnetic resonance imaging (MRI), 14-15
of orange-winged Amazon parrot, 102f-103f	Mallard duck (Anas platyrhynchos), 203f-226f
of pigeon, 200f	coelom, CT transverse view of, 225f-226f
of excretory urogram	excretory urogram, laterolateral view of, 219f,
of mallard duck, 219f, 220f	220f
of orange-winged Amazon parrot, 106f-107f	gastrointestinal positive contrast study
of gastrointestinal double contrast study	laterolateral view of, 216f
of orange-winged Amazon parrot, 98f-99f	ventrodorsal view of, 217 <i>f</i> -218 <i>f</i>
of Senegal parrot, 62f-63f	head
of gastrointestinal positive contrast study	CT transverse view of, 221f-224f
of blue and gold macaw, 136f-138f	laterolateral view of, 206f
of great horned owl, 262f-264	ventrodorsal view of, 207f
of mallard duck, 216f	pelvic limb
of Moluccan cockatoo, 174f	caudocranial view of, 213f
of orange-winged Amazon parrot, 91f-94f of peach-faced lovebird, 32f-34f	dorsoplantar view of, 215f mediolateral view of, 212f, 214f
of pigeon, 184f-186f of red tailed hawk, 239f-241f	skeleton, whole body laterolateral view, 208 <i>f</i>
of head	ventrodorsal view, 209f
of blue and gold macaw, 128f	viscera of coelom
of great horned owl, 252f	laterolateral view, 204f
of mallard duck, 206f	ventrodorsal view, 205 <i>f</i>
of Moluccan cockatoo, 164 <i>f</i>	wing
of orange-winged Amazon parrot, 80f	caudocranial view, 211f
of pigeon, 182f	mediolateral view, 210f
of red tailed hawk, 230f	Mandible
infraorbital sinus contrast study of African grey parrot, 72f	See Head.
positioning techniques	Masking tape. See Positioning techniques.

Mediolateral view	Orango winged Ameron perrot (Amerona ameronica)
of pelvic limb	Orange-winged Amazon parrot (Amazona amazonica) (Continued)
of blue and gold macaw, 134f	head
of great horned owl, 258f, 260f	laterolateral view of, 80f
of mallard duck, 212 <i>f</i> , 214 <i>f</i>	oblique view, 81f
of Moluccan cockatoo, 170f, 172f	ventrodorsal view, 82f-83f
of orange-winged Amazon parrot, 88f, 90f	liver ultrasound, sagittal view, 124f
of pigeon, 188f, 190f	MRI of head
of red tailed hawk, 235f, 237f	sagittal view, 118f-120f
positioning techniques	transverse view, 121f-123f
of foot, 7-8, 13 <i>f</i> , 14 <i>f</i>	pelvic limb
of leg, 7, 11f	craniocaudal view of, 89f
of wing, 6, 9f	dorsoplantar view of, 91f
of wing	mediolateral view of, 88f, 90f
of blue and gold macaw, 132 <i>f</i> of great horned owl, 256 <i>f</i>	skeleton, whole body laterolateral view of, 84 <i>f</i>
of mallard duck, 210f	ventrodorsal view of, 85 <i>f</i>
of Moluccan cockatoo, 168f	viscera of coelom
of orange-winged Amazon parrot, 86f	laterolateral view of, 76f, 78f
of pigeon, 186f	ventrodorsal view of, 77f, 79f
of red tailed hawk, 234f	wing
Melopsittacus undulates. See Budgerigar.	caudocranial view of, 87f
Metallic markers, in positioning, 5-7	mediolateral view of, 86f
Metals in MRI studies, 15	Orbit. See Head.
Moluccan cockatoo (Cacatua moluccensis), 159-178	Ovary. See Viscera.
CT head, 176 <i>f</i> -178 <i>f</i>	Owl, great horned. See Great horned owl.
gastrointestinal positive contrast study	P
laterolateral view of, 174f	
ventrodorsal view of, 175 <i>f</i>	Parrot. See African grey parrot; Orange-winged Amazon
head	parrot; Senegal parrot. Patella. <i>See</i> Skeleton, whole body.
laterolateral view of, 164f	Patient positioning. See Positioning techniques.
ventrodorsal view of 165 <i>f</i> pelvic limb	Peach-faced lovebird (Agapornis roseicollis), 29-37
caudocranial view of, 171 <i>f</i>	gastrointestinal positive contrast
dorsoplantar limb of, 173f	laterolateral view of, 32f-34f
mediolateral view, 170f, 172f	ventrodorsal view of, 35 <i>f</i> -37 <i>f</i>
skeleton, whole body	viscera of coelom
laterolateral view, 166f	laterolateral view of, 30f
ventrodorsal view, 167f	ventrodorsal view of, 31f
viscera of coelom	Pelleted food, 3, 9
laterolateral view, 160f, 162f	Pelvic extremity. See Foot.
ventrodorsal view, 161f, 163f	Pelvic limb
wing	See also Leg.
caudocranial view, 169f	caudocranial view
mediolateral view, 168f	of great horned owl, 259 <i>f</i> of mallard duck, 213 <i>f</i>
MRI of head, of orange-winged Amazon parrot	of Moluccan cockatoo, 171f
sagittal view, 118f-120f transverse view, 121f-123f	of orange-winged Amazon parrot, 89f
Muscle movements, 3	of pigeon, 189f
madere movements, o	of red tailed hawk, 236f
N	dorsoplantar view
Novopaque. See Barium sulfate.	of great horned owl, 261f
Nymphicus hollandicus. See Cockatiel.	of mallard duck, 215
0	of Moluccan cockatoo, 173f
	of orange-winged Amazon parrot, 91f
Object film distance (OFD), 6, 7	of pigeon, 191f
Oblique view of head, of orange-winged Amazon parrot, 81f	of red tailed hawk, 238f
Orange-winged Amazon parrot (Amazona amazonica), 75-124 cloacogram	mediolateral view
laterolateral view of, 102 <i>f</i> -103 <i>f</i>	of blue and gold macaw, 134 <i>f</i> of great horned owl, 258 <i>f</i> , 260 <i>f</i>
ventrodorsal view of, 104f-105f	of mallard duck, 212f, 214f
CT, transverse view	of Moluccan cockatoo, 170f, 172f
of coelom, 113 <i>f</i> -115 <i>f</i> , 116 <i>f</i> -117 <i>f</i>	of orange-winged Amazon parrot, 88f, 90f
head, 110 <i>f</i> -112 <i>f</i>	of pigeon, 188f, 190f
excretory urogram	of red tailed hawk, 235f, 237f
laterolateral view of, 106f-107f	Phalanges
ventrodorsal view of, 108f-109f	See also Pelvic limb; Wing.
gastrointestinal double contrast study	positioning of, 8
laterolateral view of, 98f-99f	Pharynx. See Head.
ventrodorsal view of, 100 <i>f</i> -101 <i>f</i>	Pigeon (Columba livia), 179-201
gastrointestinal positive contrast study	cloaca
laterolateral view of, 92 <i>f</i> -94 <i>f</i> ventrodorsal view of, 95 <i>f</i> -97 <i>f</i>	laterolateral view of, 200 <i>f</i>
ventiouoisai view oi, 33j-3/j	ventrodorsal view of, 201 <i>f</i>

Pigeon (Columba livia) (Continued) gastrointestinal positive contrast study laterolateral view of, 184f-186f ventrodorsal view of, 197f-199f head laterolateral view of, 182f ventrodorsal view of, 183f	References, anatomic, 2 Regurgitation of contrast medium, 11 during radiography, 3-4 Respiratory movements, 3 Respiratory series, laterolateral view of pigeon, 192f-193f Ribs. See Skeleton, whole body.
pelvic limb caudocranial view of, 189f dorsoplantar view of, 191f mediolateral view of, 188f, 190f	S Sagittal view of liver ultrasound, of orange-winged Amazon parrot,
respiratory series, laterolateral view of, 192 <i>f</i> -193 <i>f</i> skeleton, whole body laterolateral view of, 184 <i>f</i>	124f MRI head, of orange-winged Amazon parrot, 118f-120f Scapula. See Skeleton, whole body.
ventrodorsal view of, 185f viscera of coelom laterolateral view of 180f	Scatter radiation, 2 Screens, intensifying, 2-3 Sedation for radiography, 4 Senegal parrot (<i>Poicephalus senegalus</i>), 51-64
ventrodorsal view of, 181f wing caudocranial view of, 187f mediolateral view of, 186f	gastrointestinal double contrast study, laterolateral view, 62f-63f gastrointestinal positive contrast study
Poicephalus senegalus. See Senegal parrot. Positioning techniques, 4-8 for caudocranial study of thoracic extremity, 6-7, 10f	laterolateral view of, 54 <i>f</i> -57 <i>f</i> ventrodorsal view of, 58 <i>f</i> -61 <i>f</i> viscera of coelom
for craniocaudal study of pelvic limb, 7, 12 <i>f</i> devices for, 4, 4 <i>f</i> for laterolateral and ventrodorsal studies of head, 4, 5 <i>f</i> , 6 <i>f</i>	laterolateral view of, 52f ventrodorsal view of, 53f Skeleton, whole body
for laterolateral study of coelom, 4-5, 7f for mediolateral and dorsoplantar study of foot, 7-8, 13f, 14f for mediolateral study of pelvic limb, 7, 11f	lateral view of, 22f laterolateral view of African grey parrot, 70f of blue and gold macaw, 130f
of thoracic extremity, 6, 9f for ventrodorsal study of coelom, 6, 8f Proventriculus	of great horned owl, 254f of mallard duck, 208f of Moluccan cockatoo, 166f
See also Viscera. emptying of, 8 Psittacus erithacus erithacus. See African grey parrot. Public See Skeleton, whole body.	of orange-winged Amazon parrot, 83f-84f of pigeon, 184f of red tailed hawk, 232f ventral view of, 23f
Pubis. See Skeleton, whole body. Pygostyle. See Skeleton, whole body. R	ventrodorsal view of African grey parrot, 71f of blue and gold macaw, 130f
Radiology	of great horned owl, 255
equipment for	of mallard duck, 209f of Moluccan cockatoo, 167f
digital systems, 3 film-intensifying screens, 2-3, 2t	of orange-winged Amazon parrot, 85f
radiographic units, 2	of pigeon, 185f
examination for	of red tailed hawk, 233f
anesthesia, 3-4	Skull
patient preparation, 3	CT and MRI studies, 15 lateral view of, 20f
positioning devices, 2f, 4 timing of exposure, 3	ventral view of, 21f
Radius. See Wing.	Spinal cord. See Head.
Red tailed hawk (Buteo jamaicensis), 227-248	Spleen. See Viscera.
coelom, CT transverse view of, 245f-248f	Sponge, radiolucent, in positioning, 5
gastrointestinal positive contrast study laterolateral view of, 239f-241f	Sternum. See Skeleton, whole body. Sulphur-crested cockatoo (Cacatua galerita),
ventrodorsal view of, 242f-244f	151-153
head	CT head transverse view, 152, 153
laterolateral view of, 230f	Sun conure (Aratinga solstitialis), 47-49
ventrodorsal view of, 231f	viscera of coelom
pelvic limb caudocranial view of, 236f	laterolateral view, 48f ventrodorsal view, 49f
dorsoplantar view of, 238f	Survey radiography, 9, 14
mediolateral view of, 235f, 237f	Synsacrum. See Skeleton, whole body.
skeleton, whole body	Syrinx. See Viscera.
laterolateral view of, 232f	Т
ventrodorsal view of, 233f	_
viscera of coelom	Temperature of contrast medium, 9 Testes. <i>See</i> Viscera.
laterolateral view of, 228f ventrodorsal view of, 229f	Textbook references, 2
wing	Thoracic extremity. <i>See</i> Wing.
caudocranial view of, 235f	Timing of digestive tract radiography, 12
mediolateral view of, 234f	Tongue. See Head.

Trachea	Ventradorcal view (Continued)
See also Viscera.	Ventrodorsal view (Continued) of skeleton, whole body
aspiration of contrast medium, 11	of African grey parrot, 71f
Transit times in digestive tract, 12	of blue and gold macaw, 131f
Transverse view	of great horned owl, 255f
CT	of mallard duck, 209f
of coelom	of Moluccan cockatoo, 167f
of mallard duck, 225f-226f	of orange-winged Amazon parrot, 85f
of orange-winged Amazon parrot, 113f-115f,	of pigeon, 185f
116 <i>f</i> -117 <i>f</i>	of red tailed hawk, 233f
of red tailed hawk, 245 <i>f</i> -248 <i>f</i>	of viscera of coelom
of head	of African grey parrot, 69f
of blue and gold macaw, 142 <i>f</i> -145 <i>f</i>	of blue and gold macaw, 127f
of great horned owl, 268f-270f	of budgerigar, 27f
of mallard duck, 221 <i>f</i> -224 <i>f</i>	of cockatiel, 41f
of Moluccan cockatoo, 176f-178f	of Goffin cockatoo, 149f
of orange-winged Amazon parrot, 110f-112f	of great horned owl, 251f
of sulphur-crested cockatoo, 152 <i>f</i> , 153 <i>f</i>	of mallard duck, 205f
MRI of head, of orange-winged Amazon parrot, 121f-123f	of Moluccan cockatoo, 161f, 163f
Tubes, X-ray, 2	of orange-winged Amazon parrot, 77f, 79f
U	of peach-faced lovebird, 31f
Ulna. See Wing.	of pigeon, 181f of red tailed hawk, 229f
Ultra Detail Plus radiographic film, 2	of Senegal parrot, 53f
Ultrasonography of liver, of orange-winged Amazon parrot,	of sun conure, 49f
124f	of umbrella cockatoo, 157f
Umbrella cockatoo (Cacatua [Cacatua] alba), 155f-157f	Vertebra. See Skeleton, whole body.
viscera of coelom	Viscera of coelom
laterolateral view, 156f	drawings
ventrodorsal view, 157 f	of lateral view, 18f
Units, radiographic, 2	of ventral view, 19f
Urinary tract	laterolateral view
See also Excretory urogram.	of African grey parrot, 68f
radiographic contrast studies of, 14	of blue and gold macaw, 126f
V	of budgerigar, 26f
Veins. See Viscera.	of cockatiel, 40f
Ventilation, positive pressure, 3	of Goffin cockatoo, 148f
Ventral view	of great horned owl, 250f
of skeleton, 23f	of mallard duck, 204f of Moluccan cockatoo, 160f, 162f
of skull, 21f	of orange-winged Amazon parrot, 76f, 78f
of viscera, 19f	of peach-faced lovebird, 30f
Ventriculus. See Viscera.	of pigeon, 180f
Ventrodorsal view	of red tailed hawk, 228f
of cloaca	of Senegal parrot, 52f
of orange-winged Amazon parrot, 104f-105f	of sun conure, 48f
of pigeon, 201f	of umbrella cockatoo, 156
of excretory urogram, of orange-winged Amazon parrot,	ventrodorsal view
108f-109f	of African grey parrot, 69f
of gastrointestinal double contrast study, of orange-winged	of blue and gold macaw, 127f
Amazon parrot, 100f-101f of gastrointestinal positive contrast study	of budgerigar, 27f
of blue and gold macaw, 139f-141f	of cockatiel, 31f
of cockatiel, 42f-45f	of Goffin cockatoo, 149f
of great horned owl, 265f-267f	of great horned owl, 251f
of mallard duck, 217 <i>f</i> -218 <i>f</i>	of mallard duck, 205 <i>f</i> of Moluccan cockatoo, 161 <i>f</i> , 163 <i>f</i>
of Moluccan cockatoo, 175 <i>f</i>	of orange-winged Amazon parrot, 77f, 79f
of orange-winged Amazon parrot, 95f-97f	of peach-faced lovebird, 31f
of peach-faced lovebird, 35f-37f	of pigeon, 181f
of pigeon, 197 <i>f</i> -199 <i>f</i>	of red tailed hawk, 229f
of red tailed hawk, 242 <i>f</i> -244 <i>f</i>	of Senegal parrot, 53f
of head	of sun conure, 49f
of blue and gold macaw, 129f	of umbrella cockatoo, 157f
of great horned owl, 253f	Vomiting from gut anesthetic, 11
of mallard duck, 207f	TAT
of Moluccan cockatoo, 165f	W
of orange-winged Amazon parrot, 82f	Wing
of pigeon, 183f	caudocranial study, 6-7, 10f
of red tailed hawk, 231f	of blue and gold macaw, 133f
infraorbital sinus contrast study, of African grey parrot, 73f	of great horned owl, 257f
positioning techniques of coelom, 6, 8f	of mallard duck, 211 <i>f</i>
of head, 4, 5f, 6f	of Moluccan cockatoo, 169 <i>f</i> of orange-winged Amazon parrot, 87 <i>f</i>
	or orange-winged runazon parrot, o/j

Wing (Continued)
of pigeon, 187f
of red tailed hawk, 235f
mediolateral study, 6, 9f
of blue and gold macaw, 132f
of great horned owl, 256f
of mallard duck, 210f
of Moluccan cockatoo, 168f

Wing (Continued)
of orange-winged Amazon parrot, 86f
of pigeon, 186f
of red tailed hawk, 234f

X

X-ray generator, radiographic, 2